

Accepted Manuscript

Diversity, pharmacology and synthesis of bergenin and its derivatives: Potential materials for therapeutic usages

Gan B. Bajracharya

PII: S0367-326X(15)00003-9
DOI: doi: [10.1016/j.fitote.2015.01.001](https://doi.org/10.1016/j.fitote.2015.01.001)
Reference: FITOTE 3099

To appear in: *Fitoterapia*

Received date: 16 October 2014
Revised date: 1 January 2015
Accepted date: 6 January 2015



Please cite this article as: Bajracharya Gan B., Diversity, pharmacology and synthesis of bergenin and its derivatives: Potential materials for therapeutic usages, *Fitoterapia* (2015), doi: [10.1016/j.fitote.2015.01.001](https://doi.org/10.1016/j.fitote.2015.01.001)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Review

Diversity, pharmacology and synthesis of bergenin and its derivatives: Potential materials for therapeutic usages

Gan B. Bajracharya*

Faculty of Science, Nepal Academy of Science and Technology (NAST), Khumaltar, Lalitpur, Nepal

*Corresponding author's email: ganbajracharya@yahoo.com

Abstract

Bergenin, a natural secondary metabolite, has been isolated from different parts of a number of plants. It is one of active ingredients in herbal and Ayurvedic formulations. It exhibits antiviral, antifungal, antitussive, antiplasmodial, antiinflammatory, antihepatotoxic, antiarrhythmic, antitumor, antiulcerogenic, antidiabetic and wound healing properties. It has been analyzed and estimated in different plant extracts, blood and drug samples using chromatographic techniques, and pharmacokinetic studies have been made. Several bergenin derivatives were isolated and/or synthesized and were found to possess pharmacological activities. Total synthesis of bergenin and its derivatives were reported. This review article covers literature on bergenin and its derivatives until 2013. Ethnomedicinal value of bergenin containing plant materials is also highlighted. This comprehensive review provides information on the potentiality of bergenin and its derivatives for therapeutic usages.

Keywords: Analytical technique; Bergenin; Natural occurrence; Pharmacological activity; Synthesis

Contents

1. Introduction of bergenin
 2. Occurrence of bergenin in nature
 3. Ethnomedicinal values of bergenin
 4. Pharmacology on bergenin
 5. Analysis and estimation of bergenin
 6. Naturally occurred bergenin derivatives
 7. Derivatization of bergenin
 8. Total synthesis
 9. Conclusion
- Conflicts of interest
Acknowledgements
References

Abbreviations: ABTS, 2,2'-azinobis-3-ethylbenzotiazoline-6-sulfonic acid; DCC, dicyclohexylcarbodiimide; DEAD, diethyl azodicarboxylate; DIAD, diisopropyl azodicarboxylate; DMAP, N,N-dimethylamino pyridine; DME, 1,2-dimethoxyethane; DMF, N,N-dimethylformamide; DPPH, 2,2-diphenyl-1-picrylhydrazyl; EDC, ethyldimethylaminopropylcarbodiimide; IFN, interferon; IL, interleukin; MIC, minimum inhibitory concentration; PPTS, pyridinium p-toluenesulfonate; TBAF, tetra-n-butylammonium fluoride; TEMPO, 2,2,6,6-tetramethyl-1-piperidinyloxy free radical; Th, helper T cell; THF, tetrahydrofuran; TNF, tumor necrosis factor.

1. Introduction of bergenin

Natural polyphenols of plant origin are often associated with medicinal values and are often used as intermediates for industrial products and pharmacological applications. Recent studies have proved that bergenin (**1**) (Fig. 1) possesses good pharmacological properties with low side effects and little toxicity. As indicated from the result of the ^{14}C -glucose incorporation experiment in *Saxifraga stolonifera* leaves, gallic acid is considered as the glucosyl acceptor for the biosynthesis of bergenin in nature [1]. Under continuous light, bergenin present in the young leaves showed the highest incorporation of label from ^{14}C -glucose and the addition of unlabeled gallic acid enhanced the incorporation label.

Bergenin (**1**) (also known as ardisic acid B, bergenit, bergenitol, cuscutin, peltophorin and vakerin) is a C-glucoside of 4-O-methyl gallic acid (2 β -D-glucopyranosyl 4-O-methyl gallic acid δ lactone). It is an isocoumarin, hydrolysable tannin that obtain as a colorless crystal from MeOH. It has poor solubility in water, easily degrades in basic solution and its stability mostly depends on the storage conditions [2]. The initial structures of bergenin given by Tschitschibabin et al. (structure **I**) in 1928 [3] and Shimokôriyama (structure **II**) in 1950 [4] were revised by Hay et al. [5], Posternak et al. [6] and Fujise et al. [7] (Fig. 1). The structure of bergenin (**1**) was unequivocally confirmed through X-ray analysis of its 3,4,8,10,11-penta acetate derivative [8] and monohydrate [9,10].




Fig. 1. Structure of bergenin.

2. Occurrence of bergenin in nature

According to citation in the Merck Index, bergenin (**1**) was first isolated from the rhizomes of *Saxifraga (Bergenia) siberica* [11]. Later, the compound was isolated from a number of plant sources (Table 1).

Table 1

List of plants used for the isolation of bergenin (**1**).

Plant species	Family	Parts used	Yield (%)	Reference
<i>Arctostaphylos uva-ursi</i>	Ericaceae	Leafy shoot	0.0303	[12]
<i>Ardisia colorata</i>	Myrsinaceae	Fruit	0.0043	[13]
<i>Ardisia creanta</i>	Myrsinaceae	Root	2.3250	[14-16]
<i>Ardisia elliptica</i>	Myrsinaceae	Unknown	-	[17]
<i>Ardisia gigantifolia</i>	Myrsinaceae	Rhizome	0.0148	[18]
<i>Ardisia japonica</i>	Myrsinaceae	Aerial	0.0225	[19,20]
<i>Ardisia punctata</i>	Myrsinaceae	Root	-	[21]
<i>Arisaema franchetianum</i>	Araceae	Tuber	0.0001	[22]
<i>Astilbe chinensis</i>	Saxifragaceae	Rhizome	0.0397	[23,10,24]
<i>Astilbe myriantha</i>	Saxifragaceae	Unknown	-	[25]
<i>Astilbe rivularis</i>	Saxifragaceae	Rhizome	0.0002	[26,27]

<i>Astilbe thunbergii</i>	Saxifragaceae	Rhizome	0.4953	[28,29]
<i>Bergenia cordifolia</i>	Saxifragaceae	Leaf	-	[30-32]
<i>Bergenia cordifolia</i>	Saxifragaceae	Rhizome	0.5688	[33]
<i>Bergenia crassifolia</i>	Saxifragaceae	Rhizome	3.3632	[34,3,5]
<i>Bergenia crassifolia</i>	Saxifragaceae	Leaf	-	[35]
<i>Bergenia ligulata (ciliata)</i>	Saxifragaceae	Rhizome	0.0760	[36-42]
<i>Bergenia purpurascens</i>	Saxifragaceae	Rhizome	1.5935	[43,44]
<i>Bergenia scopulosa</i>	Saxifragaceae	Rhizome	-	[45,46]
<i>Bergenia stracheyi</i>	Saxifragaceae	Whole plant	-	[47,48]
<i>Bergenia stracheyi</i>	Saxifragaceae	Rhizome	2.0425	[49,50]
<i>Brachystemma calycinum</i>	Caryophyllaceae	Aerial	0.0002	[51]
<i>Caesalpinia decapetala</i>	Fabaceae	Root	0.0001	[52]
<i>Caesalpinia digyna</i>	Fabaceae	Root	0.0650	[53-57]
<i>Caesalpinia mimosoides</i>	Fabaceae	Root	0.0088	[58]
<i>Cenostigma gardnerianum</i>	Leguminosae	Stem bark	0.0747	[59]
<i>Cenostigma macrophyllum</i>	Leguminosae	Stem bark	0.0747	[60]
<i>Cimicifuga foetida</i>	Ranunculaceae	Rhizome	0.0125	[61,62]
<i>Connarus monocarpus</i>	Connaraceae	Root	-	[63]
<i>Corylopsis coreana</i>	Hamamelidaceae	Leaf	0.8611	[64]
<i>Corylopsis spicata</i>	Hamamelidaceae	Bark	-	[3,65]
<i>Corylopsis willmottiae</i>	Hamamelidaceae	Whole plant	0.0040	[66]
<i>Diospyros Sanja-Minika</i>	Ebenaceae	Wood	3.6402	[67]
<i>Dipterocarpus grandiflorus</i>	Dipterocarpaceae	Stem	0.1600	[68]
<i>Dryobalanops aromatica</i>	Dipterocarpaceae	Stem bark	0.0025	[69,70]
<i>Dryobalanops sp.</i>	Dipterocarpaceae	Heartwood	-	[71]
<i>Endopleura uchi</i>	Humiriaceae	Bark	1.1347	[72-74]
<i>Endopleura uchi</i>	Humiriaceae	Fruit	-	[75]
<i>Ficus racemosa</i>	Moraceae	Bark	-	[76-79]
<i>Flueggea microcarpa</i>	Euphorbiaceae	Leaf	-	[80-82]
<i>Fluggea virosa</i>	Euphorbiaceae	Aerial	-	[83-85]
<i>Garcinia malaccensis</i>	Clusiaceae	Stem bark	0.0012	[86]
<i>Gendarussa vulgaris</i>	Acanthaceae	Aerial	0.0003	[87]
<i>Glochidion obliquum</i>	Euphorbiaceae	Leaf	0.0023	[88]
<i>Glochidion obovatum</i>	Euphorbiaceae	Leaf	0.0006	[89]
<i>Hopea sangal</i>	Dipterocarpaceae	Stem bark	0.0017	[90]
<i>Hopea utilis</i>	Dipterocarpaceae	Stem wood	0.1800	[91]
<i>Humiria balsamifera</i>	Humiriaceae	Aerial	-	[9,92]
<i>Macaranga peltata</i>	Euphorbiaceae	Bark	0.4500	[93]
<i>Mallotus anisopodus</i>	Euphorbiaceae	Aerial	0.0002	[94]
<i>Mallotus japonicus</i>	Euphorbiaceae	Bark	0.8796	[95-99]
<i>Mallotus japonicus</i>	Euphorbiaceae	Cortex	1.9500	[100-102]
<i>Mallotus philippinensis</i>	Euphorbiaceae	Leaf	0.0001	[103]
<i>Mallotus philippinensis</i>	Euphorbiaceae	Stem bark	0.6500	[104-107]
<i>Mallotus repandus</i>	Euphorbiaceae	Stem	0.0380	[108,109]
<i>Mallotus roxburghianus</i>	Euphorbiaceae	Leaf	0.0075	[110]
<i>Peltiphyllum peltatum</i>	Saxifragaceae	Rhizome	0.0092	[111]
<i>Peltoboykinia watanabei</i>	Saxifragaceae	Rhizome	-	[112]
<i>Peltophorum africanum</i>	Fabaceae	Stem bark	1.2000	[113-116]
<i>Peltophorum africanum</i>	Fabaceae	Root	2.0000	[116]
<i>Peltophorum ferruginum</i>	Fabaceae	Bark	-	[117]
<i>Peltophorum inerme</i>	Fabaceae	Flower	-	[118]
<i>Peltophorum pterocarpum</i>	Fabaceae	Flower	0.1008	[119,120]

<i>Peltophorum pterocarpum</i>	Fabaceae	Wood	0.0211	[121]
<i>Pentaclethra macrophylla</i>	Mimosaceae	Root	0.0004	[122]
<i>Phyllanthus columnaris</i>	Euphorbiaceae	Root bark	0.0014	[123]
<i>Phyllanthus flexuosus</i>	Euphorbiaceae	Stem bark	-	[124]
<i>Phyllanthus wightianus</i>	Euphorbiaceae	Whole plant	-	[125]
<i>Pulicaria wightiana</i>	Compositae	Aerial	0.0007	[126]
<i>Pulsatilla koreana</i>	Ranunculaceae	Root	0.0001	[127]
<i>Rivea hypocrateriformis</i>	Convolvulaceae	Stem	0.0070	[128]
<i>Rodgersia aesculifolia</i>	Saxifragaceae	Rhizome	-	[129]
<i>Rodgersia pinnata</i>	Saxifragaceae	Unknown	0.0001	[130]
<i>Rodgersia sambucifolia</i>	Saxifragaceae	Root	3.1250	[16]
<i>Sacoglottis gabonensis</i>	Humiriaceae	Bark	0.0500	[131-135]
<i>Sacoglottis uchi</i>	Humireaceae	Bark	0.1000	[136]
<i>Saxifraga melanocentra</i>	Saxifragaceae	Aerial	-	[137]
<i>Saxifraga stolonifera</i>	Saxifragaceae	Aerial	-	[138-141]
<i>Securinega melanthesoides</i>	Euphorbiaceae	Leaf	-	[142]
<i>Securinega virosa</i>	Euphorbiaceae	Leaf	0.0122	[143]
<i>Shorea leprosula</i>	Dipterocarpaceae	Heartwood	-	[144]
<i>Shorea robusta</i>	Dipterocarpaceae	Leaf	0.0750	[145]
<i>Shorea robusta</i>	Dipterocarpaceae	Root	-	[146]
<i>Streptocaulon griffithii</i>	Asclepiadaceae	Root	0.0022	[147]
<i>Teramnus labialis</i>	Fabaceae	Aerial	-	[148]
<i>Tridax procumbens</i>	Heliantheae	Aerial	0.0017	[149]
<i>Tripterospermum chinense</i>	Gentianaceae	Aerial	-	[150]
<i>Vateria indica</i>	Dipterocarpaceae	Leaf	0.2600	[151,152]
<i>Vateria indica</i>	Dipterocarpaceae	Seed	-	[153]
<i>Vateria indica</i>	Dipterocarpaceae	Stem bark	1.7647	[154]
<i>Vatica albiramis</i>	Dipterocarpaceae	Stem	0.0833	[155]
<i>Vatica bantamensis</i>	Dipterocarpaceae	Leaf	0.1100	[156]
<i>Vatica diospyroides</i>	Dipterocarpaceae	Stem	0.0640	[157]
<i>Vatica mangachpoi</i>	Dipterocarpaceae	Leaf	-	[158,159]
<i>Vatica pauciflora</i>	Dipterocarpaceae	Stem bark	0.0840	[160]
<i>Viburnum nervosum</i>	Caprifoliaceae	Root	-	[161,162]
<i>Woodfordia fruticosa</i>	Lythraceae	Stem	-	[163]

Yield percentage is calculated in dried plant material weight basis and the highest isolated yield is recorded from the available data.

As can be seen from Table 1, rhizomes of *Ardisia creanata*, *Bergenia crassifolia*, *Bergenia purpurascens*, *Bergenia stracheyi*, *Peltophorum africanum* and *Rodgersia sambucifolia* are the major sources of bergenin (1). Barks of *Diospyros Sanja-Minika*, *Endopleura uchi*, *Mallotus japonicus*, *Peltophorum africanum* and *Vatica indica* also contain a high amount of the compound. Leaf of *Corylopsis coreana* is another important source of bergenin. When deposition of bergenin is high in the plant material, simple extraction with acetone, methanol or water followed by concentration and crystallization could be efficient for its isolation [5,97,124,154]. Otherwise, Soxhlet extraction followed by silica gel column chromatography is often employed for the isolation of bergenin. However, the conventional method alone is not efficient to isolate bergenin since repeated silica gel column chromatography, the use of expensive reversed phase adsorbents, preparative high performance liquid chromatography (HPLC) and/or recrystallizations are often essential [19,43,49,64,101,128]. A rapid extraction and purification of bergenin by microwave-assisted extraction coupled with high-speed counter-current chromatography (HSCCC) was reported by Deng et al., in which sample was extracted with 60% aqueous methanol with solvent/sample ratio of 10/1

(mL/g) at 60 °C for 15 min followed by direct HSCCC purification [16]. This method was found efficient but limited to small scale. Recently, we have reported an efficient, simple methodology for purification of bergenin (**1**) from *Bergenia purpurascens* rhizome extract through column chromatography using alumina as an adsorbent [44]. Utilizing this protocol, bergenin can be isolated exclusively. MAE/HSCCC

3. Ethnomedicinal values of bergenin

Bergenin (and its congeners) occur in a large number of plants and is considered as an active ingredient in the plant extracts. Bergenin containing herbs have been used as a folk medicine in Asia (India, China including Nepal) since at least the 7th century [164-168]. It is one of the active pharmaceutical ingredients of Ayurvedic herbal drugs and formulations [169]. In Ayurvedic formulations, bergenin containing plant materials such as *Bergenia ligulata* is used in lithiasis, dysuria and polyuria; *Caesalpinia digyna* is used as astringent and antipyretic; *Peltophorum pterocarpum* is used for dysentery and muscular pains; *Vateria indica* is considered to be effective against bronchitis, gonorrhea and syphilis; *Woodfordia fruticosa* is used as tonic and sedative; etc. Reviews on the ethnomedicinal values of bergenin containing plant materials have highlighted the importance of bergenin [2,164,170-179].

4. Pharmacology on bergenin

4.1. Evaluation in infectious diseases

Antibacterial activity

Bergenin (**1**) was found ineffective against *Escherichia coli*, *Salmonella enteritidis*, *Shigella sonnei*, *Pseudomonas aeruginosa*, *Serratia marcescens*, *Klebsiella pneumoniae*, *Salmonella paratyphi*, *Salmonella typhi*, *Salmonella typhimurium*, *Shigella flexneri*, *Proteus vulgaris*, *Enterococcus faecalis*, *Staphylococcus aureus*, *Bacillus subtilis* and *Erwinia* sp. [44,50,72,120]. Inhibitory effect of bergenin on *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Beta streptococcus* and *Aeruginosus bacillus* was reported [180]. Antitussive effect of bergenin has been reported elsewhere [181,182].

Antiviral activity

Piacente et al. reported a weak anti-HIV-1 activity of bergenin (**1**) in C8166 cells infected with HIV-1_{MN} (X4 virus) with an effective concentration (EC₅₀) value of 40 µg/mL. And, bergenin inhibited the binding of GP120 to sCD4 in a dose-dependent manner [19]. On NS3 serine protease activity assay, bergenin has displayed a weak activity against Hepatitis C virus (IC₅₀ = 1.71 mM) [137]. Antiviral activity of bergenin against herpes simplex virus type-1 showed an IC₅₀ value of <6.25 µg/mL [27]. Bergenin had no inhibitory property against HIV-1 reverse transcriptase and integrase [114].

Antifungal activity

Bergenin (**1**) displayed antifungal activity against *Candida albicans*, *Candida tropicalis* and *Candida guilliermondii* with MIC values of 14.9, 14.9 and 29.8 µM, respectively, while a lower activity was observed against filamentous fungi *Aspergillus flavus*, *Aspergillus nidulans* and *Aspergillus niger* [72]. Antifungal activity of bergenin against *Trichophyton mentagrophytes* (MIC 250 µg/mL), *Epidermophyton floccosum* (MIC 500 µg/mL), *Trichophyton rubrum* (MIC 500 µg/mL), *Aspergillus niger* (MIC 500 µg/mL) and *Botrytis cinerea* (MIC 250 µg/mL) was reported by Raj et al. [120]. The monosodium salt of bergenin, obtained by treating bergenin with one molar equivalent of NaOH in water, was found effective against *Alternaria alternata*, *Alternaria brassicae*, *Alternaria carthami*, *Fusarium udum*, *Fusarium oxysporum* f. sp. *ciceri*, *Curvularia lunata* and *Erysiphe pisi* [183].

Wound healing effect

Effective dose (ED₅₀) of bergenin (**1**) for burn wound healing was found to be 190 µg/wound in mice [29]. Wound healing effect of bergenin was also reported by Mukherjee et al. [145].

4.2. Evaluation as antiparasitics and insecticides**Antiplasmodial activity**

Bergenin (**1**) displayed a good activity with IC₅₀ value of 2.4 µg/mL against chloroquine sensitive strain of *Plasmodium falciparum* (D10) indicating it is a new potential antimalarial agent [42].

Antifeedent activity

Bergenin (**1**) exhibited significant antifeedant activity against lepidopterous insects [26].

Trypanocidal activity

Bergenin (**1**) showed an inhibitory effect on the growth of *Trypanosoma brucei* with an IC₅₀ value of 1 mM [84].

4.3. Evaluation in immunological and inflammatory diseases**Antiinflammatory activity**

Swarnalakshmi et al. have reported that bergenin (**1**) produces a dose dependent inhibition of carrageenin induced rat paw oedema [119]. Nazir et al. have reported antiarthritic activity of bergenin in adjuvant-induced arthritic balb/c mice [49]. A flow cytometric study revealed that bergenin inhibits the production of proinflammatory Th1 cytokines (IL-2, IFN-γ and TNF-α) and promotes the production of Th2 cytokines (IL-4 and IL-5). De Oliveira et al. have reported antinociceptive and antiinflammatory properties of bergenin owing to the inhibition of IL-1β and TNF-α release [59]. Nunomura et al. reported antiinflammatory effect of bergenin against cyclooxygenase 1 (Inhibition concentration (IC₅₀) = 107.2 µM), cyclooxygenase 2 (IC₅₀ = 1.2 µM) and phospholipase A₂ (IC₅₀ = 156.6 µM) [73]. Jachak et al. have reported that bergenin was an inhibitory principle in the ethyl acetate extract of *Tridax procumbens* for antiinflammatory effect [149]. Antiinflammatory activity of bergenin against 12-O-tetradecanoylphorbol-13-acetate (TPA)-induced inflammation in mice was evaluated by Zhang et al. [121]. Bergenin displayed a low percent inhibitory ratio (34 ± 4.7% at 0.5 mg/ear) compared to indomethacin (96 ± 4.7% at 1.0 mg/ear).

Antihepatotoxic activity

The hepatoprotective effect of bergenin (**1**) utilizing CCl₄-induced rat hepatocytes was initially reported by Hikino et al. [184]. Kim et al. reported bergenin reduces the levels of glutamic pyruvic transaminase (GPT) and sorbitol dehydrogenase (SDH) in the culture medium released from the CCl₄-intoxicated rat hepatocytes [101]. The release of GPT and SDH was blocked 62.1% and 49.9% respectively at 100 µM concentration of bergenin. The antihepatotoxicity of bergenin was also evidenced by recovering of glutathione content (up to 25.3%), and elevating the activities of glutathione S-transferase (up to 51.5%) and glutathione reductase (up to 29.1%) at 300 µM concentration. The release of GPT and SDH was blocked by bergenin on D-galactosamine-intoxicated rat hepatocytes. At the same time, DNA synthesis was increased indicating liver protection activity of bergenin [185,186,99]. It was found that enzymatic activities of alanine/aspartate aminotransferase, sorbitol dehydrogenase, γ-glutamyltransferase, glutathione S-transferase and glutathione reductase were restored towards normalization in the experimental models after oral administration of bergenin.

Rats pretreated with bergenin significantly reduced 2,4-dinitrophenyl hydrazine-induced lipid peroxidation in the liver, brain and red blood cell [134,135].

4.4. Evaluation in blood and cardiovascular diseases

Platelet aggregation inhibition

Bergenin (**1**) inhibited platelet aggregation in human blood [86]. It showed marked inhibitory effect caused by the three inducers arachidonic acid ($IC_{50} = 240.6 \mu M$), adenosine diphosphate ($IC_{50} = 120.3 \mu M$) and collagen ($IC_{50} = 60.9 \mu M$).

Antiarrhythmic activity

Bergenin (**1**) has shown efficacy in restoring sinus rhythm in $BaCl_2$ -induced arrhythmic rats, and it has shortened the duration of ventricular premature beat, tachycardia and fibrillation after ligation and reperfusion [83]. Bergenin could increase the atria fibrillation threshold in urethane induced anesthetized rabbits indicating its potentiality to treat cardiac arrhythmias.

4.5. Evaluation in oncological diseases

Anticancer activity

Bergenin (**1**) showed a weak activity ($IC_{50} = 44 \mu M$) against the Murine Breast Cancer Cell Line, FM3A [13]. Growth inhibition of HepG2 cells was achieved [187]. Anti-proliferative effects of bergenin on human prostate cancer cell lines LNCaP and DU145 were studied [64]. The cytotoxic activity of bergenin against several cell lines was also studied by Wibowo et al. [69] and Xue et al. [24]. Bergenin has exhibited inhibitory effects against *Epstein-Barr* virus early antigen (EBV-EA) activation induced with TPA in *Raji cells* and against skin tumor promotion in mouse skin carcinogenesis [121]. It exhibited melanogenesis inhibitory activity in α -melanocyte-stimulating hormone (α -MSH)-stimulated B 16 melanoma cells therefore may be valuable as the potential skin whitening agents. Bergenin exhibited protection effect in γ -radiation induced DNA (pBR322) damage [55]. No antitumor activity of bergenin on human gastric carcinoma cell line MGC-803 was reported [52].

Antioxidant activity

Reactions of pulse radiolytically generated hydroxyl ($\bullet OH$) radicals with bergenin have been studied, which showed that bergenin radical products are formed by $\bullet OH$ radical addition to the phenyl ring and H-atom abstraction from the C ring in bergenin [188]. In most of the cases, $\bullet OH$ radical reacts with natural polyphenols to produce phenoxyl radicals, in contrast, in the case of bergenin, reducing radical adducts are the major transients formed, which may react with oxygen forming peroxy type radicals. Therefore, bergenin may not act as a potent antioxidant in preventing free radical induced oxidative damage; however, may act as a pro-oxidant and exhibit antitumor activity. Theoretical calculations on the formation of the radical derivatives of bergenin using $\bullet H$, $\bullet OH$, $\bullet CH_3$ and $\bullet CCl_3$ revealed that the methoxy group at 6-position is the most favourable site for a radical attack [136]. To confirm antioxidant activity of bergenin, β -carotene, DPPH and a heterogeneous Fenton assays were carried out.

Bergenin (**1**) itself is a good scavenger of hydroxyl radicals but not so effective in scavenging other free radicals like superoxide radical and DPPH [33,35,41,42,50,52,55,64,110,111,148,188,189]. Significant antioxidant activity of bergenin in hydrogen peroxide, ABTS, DPPH and inhibition of lipid peroxidation assays with IC_{50} values 32.54, 75.06, 165.35 and 365.12 $\mu g/mL$, respectively was reported by Srinivasan et al. [54]. Bergenin has displayed low antioxidant activity in nitric oxide

method ($IC_{50} = 785.63 \mu\text{g/mL}$) and deoxy ribose method ($IC_{50} = 815.63 \mu\text{g/mL}$), and was found to be inactive in scavenging hydroxyl radical by DNA method and superoxide radical by alkaline dimethyl sulfoxide method. DPPH radical scavenging activity of bergenin was also reported by Sumino et al. [13] and Zamarrud et al. [128].

4.6. Evaluation in alimentary tract and metabolic diseases

Gastroprotective activity

Bergenin (**1**) showed a dose dependent antisecretory effect on gastric secretion in pylorus ligated rats and antiulcerogenic activity in stressed rat ulcers [190]. The effectiveness may be due to inhibition of acetylcholine release [191]. Bergenin showed gastroprotective effect by increasing prostaglandin production [192]. It inhibited the bovine adrenal tyrosine hydroxylase activity by 29% at $20 \mu\text{g/mL}$ concentration [102].

Antidiabetic activity

Bergenin (**1**) at 10 mg/kg oral dose to streptozotocin-nicotinamide induced diabetic rats was found to reduce blood glucose level significantly in oral glucose tolerance test [56]. It reversed plasma lipid profile (total cholesterol, triglycerides and lipoproteins) to normal values, decreased lipid peroxides, and increased superoxide dismutase and catalase in liver illustrating its antidiabetic, hypolipidemic and antioxidant activities in Type 2 diabetic rats. Histopathological studies demonstrated a considerable regenerative effect on the β cells of pancreas attributing a positive effect of bergenin on the endocrine cells to produce insulin. Potentially antidiabetic activity of bergenin was reported by Li et al. since it inhibited human protein tyrosine phosphatase 1B (hPTP1B) activity with IC_{50} value of $157 \mu\text{M}$ [20].

Urease inhibitor

Bergenin (**1**) was found to inhibit the *Bacillus pasteurii* urease [105]. Urease produced by bacteria such as *Helicobacter pylori* in gastrointestinal tract catalyzes the hydrolysis of urea to produce ammonia and carbon dioxide, and permits bacteria to grow and colonize at the low pH leading gastric and peptic ulceration and associated cancer. Molecular docking study showed that bergenin penetrates into the active site of urease preventing the access to urea, thereby involving in antiulcer activity.

Lipolysis effect (antiobesity activity)

Bergenin (**1**) has enhanced norepinephrine-induced lipolysis in endogenous lipid droplets, slightly stimulated adrenocorticotrophic hormone-induced lipolysis and inhibited insulin-induced lipogenesis from glucose in fat cells obtained from rat epididymal adipose tissues [28]. Therefore, bergenin may be effective in formulations for the treatment of obesity.

Hypolipidemic activity

Oral administration of bergenin (**1**) to hyperlipidemic rats significantly decreased serum total lipid but not much change in serum cholesterol and triglycerides for 14 days [82]. However after 21 days of feeding, serum cholesterol, triglycerides, low-density lipoprotein-cholesterol levels were significantly reduced, while the serum high-density-cholesterol level was elevated.

4.7. Evaluation in renal diseases

Effect of bergenin (**1**) on urolithiasis induced by 3% glycolic acid in albino rats was evaluated; however, it exhibited less significant antiurolithiatic activity [193].

5. Analysis and estimation of bergenin

Thin layer chromatography (TLC), HPLC and Liquid chromatography-tandem mass spectrometry (LC-MS/MS) techniques have been employed for the estimation of bergenin (**1**) in the plant materials. TLC method was used for the quantification of bergenin in different parts of different *Bergenia* species [194-197,35] and *Mallotus philippensis* [198]. Several researchers have used HPLC method to determine bergenin in genus *Bergenia* [39,197,199-202], *Endopleura uchi* [73], *Peltophorum pterocarpum* [203], *Ardisia japonica* [204] and *Ficus racemosa* [205,206]. About 3.2% of bergenin was found to be deposited in the rhizomes of *Bergenia ligulata*, *Bergenia ciliata* and *Bergenia stracheyi* [39,199]. Up to 9.77% of bergenin was found to be deposited in the rhizome of *Bergenia purpurascens* distributed in Yulong reservoir of Lijiang County, China [200]. Bark of *Endopleura uchi* contained 3.19% of bergenin [73]. Flower of *Peltophorum pterocarpum* contains 0.40% of bergenin [203]. The amount of bergenin in stem bark of *Ficus racemosa* was estimated by Veerapur et al. and Ahmed and Urooj separately and found to be 0.15% and 0.89%, respectively [205,206].

Pharmacokinetic characterization of the substance is essential during drug development process. Drug absorption, distribution, metabolism and excretion are the key factors for development of effective drug. HPLC method has been employed for the determination of bergenin (**1**) in rat plasma [207,208]. After intravenous administration of bergenin in Wistar rats at the dose of 11.25 mg/kg, blood samples were collected at different intervals [207]. The study of plasma concentration-time curve of bergenin indicated that bergenin distributed widely and eliminated with moderate velocity in rat. On the other hand, when bergenin was given to rats by oral route at a single dose of 22.5 mg/kg, no satisfied plasma concentration data were obtained perhaps indicating that bergenin is easily degraded in the digestive system, including quick metabolism or a poor absorption in gastrointestinal tract. Upon intravenous administration of bergenin to the rats, bergenin was rapidly distributed in blood plasma and eliminated with the half-lives of 3 and 33 min respectively and were not related to the administration doses [208]. Shi et al. reported determination of bergenin in rat urine, feces and tissues after intravenous administration of formulated bergenin at a single dose of 22.5 mg/kg body weight of Wistar rats [209]. The study demonstrated that 35.36% and <10% of bergenin was recovered from urine and feces, respectively. Comparison to the tissues (liver, kidney, lung, heart, spleen and brain) examined, kidney accumulated the highest bergenin concentration and the longest drug resident time, while it was less significantly distributed into the brain owing to its hydrophilic property.

Yu et al. reported quantitation of bergenin and pharmacokinetic study in human plasma by LC-MS/MS [210]. The blood samples from healthy volunteers were collected at different intervals after oral administration of 250 mg of bergenin. The maximum concentration of 66.6 ng/mL occurred at 2.0 ± 0.9 h and the elimination half-time was 3.7 ± 2.4 h. HPLC-MS/MS method was used by Wang et al. to determine bergenin in human plasma [211]. This method was found to be suitable for the determination of low bergenin concentration (lowest limit of quantification = 0.25 ng/mL) in human plasma after oral administration of bergenin tablet (250 mg) and pharmacokinetic parameters (absorption half life = 0.852 h, time to peak concentration = 2.125 h, maximum plasma concentration = 15.915 ng/mL, elimination half life = 4.198 h etc.) were given. Bergenin content in *Caesalpinia digyna* roots was estimated by LC-MS and found to be contained nearly 30% in the plant extracts [55].

The interaction between bergenin and human serum albumin (HAS) in membrane mimetic environment was studied by Zhang [212]. This study indicated that bergenin bound to HAS mainly by a hydrophobic interaction.

Qin et al. reported that oral bioavailability of the drug can be increased utilizing bergenin-phospholipid complex, which was prepared by refluxing bergenin (**1**) with phospholipid (bergenin to phospholipid ratio = 0.9 w/w) in anhydrous ethanol at 60 °C for 2 h. This complex was orally

administered to rats and blood plasma was analyzed by HPLC. The study showed that the relative bioavailability was significantly increased to 439% of bergenin [213].

Voltammetric determination of bergenin was demonstrated by Chen et al. [214]. A 4-(2-pyridylazo)-resorcinol polymer film modified glassy carbon electrode was used to accumulate bergenin leading enhancement of the oxidation peak current depending on the concentration. This method was used to determine bergenin in tablets and urine samples. The electrocatalytic oxidation of bergenin was also investigated on the surface of a multi-wall carbon nanotubes modified carbon paste electrode to determine the amounts of bergenin in tablets [215].

6. Naturally occurred bergenin derivatives

A list of bergenin derivatives isolated from the plant materials is depicted in Table 2 and the structures of the compounds are shown in Fig. 2.

Table 2

Bergenin derivatives (2-35) isolated from plants.

Compound isolated	Plant species	Part used	Yield (%)	References
Norbergenin (2)	<i>Ardisia colorata</i>	Fruit	0.0170	[13]
	<i>Ardisia japonica</i>	Leaf	0.0129	[19]
	<i>Bergenia crassifolia</i>	Root	-	[36,38]
	<i>Caesalpinia digyna</i>	Root	-	[53]
	<i>Corylopsis coreana</i>	Leaf	0.0049	[64]
	<i>Corylopsis spicata</i>	Bark	-	[3]
	<i>Mallotus japonicus</i>	Bark	0.0028	[98]
	<i>Peltophorum africanum</i>	Bark	-	[113]
	<i>Saxifraga stolonifera</i>	Whole plant	-	[139]
	<i>Shorea leprosula</i>	Heartwood	-	[144]
	<i>Woodfordia fruticosa</i>	Stem	-	[163]
8-O-Methylnorbergenin (3)	<i>Saxifraga stolonifera</i>	Whole plant	0.0004	[216]
4-O-Galloylnorbergenin (4)	<i>Mallotus japonicus</i>	Bark	0.0005	[98]
11-O-Galloylnorbergenin (5)	<i>Mallotus japonicus</i>	Bark	0.0147	[97]
	<i>Mallotus japonicus</i>	Bark	0.0004	[98]
Demethoxybergenin (6)	<i>Ardisia colorata</i>	Fruit	0.0036	[13]
8,10-Di-O-methylbergenin (7)	<i>Ardisia japonica</i>	Leaf	0.0090	[19]
	<i>Macaranga peltata</i>	Heartwood	0.0133	[93]
3,8,10-Tri-O-methylbergenin (8)	<i>Macaranga peltata</i>	Heartwood	0.0033	[93]
8,10,11-Tri-O-methylbergenin (9)	<i>Macaranga peltata</i>	Heartwood	0.0008	[93]
Rivebergenin A (10)	<i>Rivea hypocrateriformis</i>	Stem	0.0035	[128]
11-O-Acetylbergenin (11)	<i>Flueggea virosa</i>	Aerial	-	[85]
	<i>Vitis repens</i>	Unknown	-	[217]
4-O-Galloylbergenin (12)	<i>Ardisia gigantifolia</i>	Rhizome	0.0001	[18]
	<i>Bergenia purpurascens</i>	Root	0.0169	[43]
	<i>Bergenia scopulosa</i>	Rhizome	-	[46]
	<i>Corylopsis willmottiae</i>	Whole plant	0.0031	[66]
	<i>Mallotus japonicus</i>	Bark	0.1730	[97]
	<i>Mallotus japonicus</i>	Bark	0.0173	[98]
	<i>Mallotus philippinensis</i>	Leaf	0.0004	[103]
	<i>Ardisia gigantifolia</i>	Rhizome	0.0002	[18]
11-O-Galloylbergenin (13)	<i>Astilbe chinensis</i>	Rhizome	0.0006	[24]
	<i>Bergenia ciliata</i>	Root	-	[218]
	<i>Bergenia ligulata</i>	Rhizome	0.0043	[42,219]
	<i>Bergenia purpurascens</i>	Root	0.1084	[43]

	<i>Caesalpinia digyna</i>	Root	0.0032	[57]
	<i>Corylopsis coreana</i>	Leaf	0.0450	[64]
	<i>Corylopsis willmottiae</i>	Whole plant	0.3188	[66]
	<i>Mallotus japonicus</i>	Bark	0.1914	[97]
	<i>Mallotus japonicus</i>	Bark	0.0070	[98]
	<i>Mallotus philippinensis</i>	Stem bark	0.0165	[220]
	<i>Peltiphyllum peltatum</i>	Rhizome	0.0007	[111]
3,4-Di-O-galloylbergenin (14)	<i>Mallotus japonicus</i>	Bark	0.0008	[98]
3,11-Di-O-galloylbergenin (15)	<i>Bergenia crassifolia</i>	Root	-	[221]
4,11-Di-O-galloylbergenin (16)	<i>Corylopsis willmottiae</i>	Whole plant	0.0004	[66]
	<i>Mallotus japonicus</i>	Bark	0.0006	[98]
3,4,11-Tri-O-galloylbergenin (17)	<i>Mallotus japonicus</i>	Bark	0.0004	[98]
Bergecin A (18)	<i>Bergenia stracheyi</i>	Whole plant	0.0001	[48]
Bergecin B (19)	<i>Bergenia stracheyi</i>	Whole plant	0.0001	[48]
10-O-(p-Hydroxybenzoyl)bergenin (20)	<i>Saxifraga melanocentra</i>	Aerial	-	[137]
11-O-(p-Hydroxybenzoyl)bergenin (21)	<i>Astilbe chinensis</i>	Rhizome	0.0002	[24]
	<i>Vatica bantamensis</i>	Leaf	0.0005	[156]
11-O-Vanilloylbergenin (22)	<i>Ardisia crenata</i>	Root	-	[15]
	<i>Vatica bantamensis</i>	Leaf	0.0007	[156]
11-O-(3'-O-methylgalloyl)bergenin (23)	<i>Ardisia gigantifolia</i>	Rhizome	0.0002	[18]
	<i>Astilbe chinensis</i>	Rhizome	0.0013	[24]
	<i>Corylopsis willmottiae</i>	Whole plant	0.0001	[66]
11-O-(4'-O-Methylgalloyl)bergenin (24)	<i>Crassula</i> cv. 'Himaturi'	Whole plant	0.0735	[222]
	<i>Saxifraga melanocentra</i>	Aerial	-	[223]
11-O-(3',4'-Di-O-methylgalloyl)bergenin (25)	<i>Ardisia crenata</i>	Root	-	[15]
11-O-Veratroylbergenin (26)	<i>Ardisia gigantifolia</i>	Rhizome	0.00015	[18]
11-O-Syringylbergenin (27)	<i>Ardisia crenata</i>	Root	-	[15]
	<i>Ardisia gigantifolia</i>	Rhizome	0.0002	[18]
	<i>Corylopsis willmottiae</i>	Whole plant	0.0005	[66]
	<i>Vatica bantamensis</i>	Leaf	0.0002	[156]
11-O-(E)-p-Coumaroylbergenin (28)	<i>Peltophorum africanum</i>	Bark	-	[113]
	<i>Vatica bantamensis</i>	Leaf	0.0002	[156]
11-O-(Z)-p-Coumaroylbergenin (29)	<i>Vatica bantamensis</i>	Leaf	0.0005	[156]
11-O-(E)-Caffeoylbergenin (30)	<i>Securinega virosa</i>	Leaf	0.0039	[143]
11-O-(E)-Ferulaylbergenin (31)	<i>Vatica bantamensis</i>	Leaf	0.0002	[156]
11-O-(Z)-Ferulaylbergenin (32)	<i>Vatica bantamensis</i>	Leaf	0.0001	[156]
11-O-(E)-Sinapoylbergenin (33)	<i>Vatica bantamensis</i>	Leaf	0.0011	[156]
Rivebergenin B (34)	<i>Rivea hypocrateriformis</i>	Stem	0.0049	[128]
Dimer of bergenin (35)	<i>Astilbe rivularis</i>	Rhizome	0.0010	[27]

Percentage of isolated yield is recorded in dried plant material weight basis from the available data.




Fig. 2. Bergenin derivatives (**2-35**) isolated from the plant materials.

Norbergenin (**2**) has shown to have antioxidant [224,13], gastroprotective [192,102], anti-HIV [19], and antiarthritic [49] activities. 11-O-Acetyl bergenin (**11**) showed antitrypanosomal activity with IC_{50} value of 0.17 mM against trypomastigotes of *Trypanosoma evansi* [217]. DPPH free-radical scavenging activity of 11-O-galloylbergenin (**13**) [111,42,18]; 4-O-galloylbergenin (**12**), 11-O-(3'-O-methylgalloyl)bergenin (**23**), 11-O-veratroylbergenin (**26**) and 11-O-syringylbergenin (**27**) [18]; rivebergenin A (**10**) and rivebergenin B (**34**) [128]; bergecin A (**18**) and bergecin B (**19**) [48]; 11-O-(4'-O-methylgalloyl)bergenin (**24**) [222]; and 11-O-(*E*)-caffeoylbergenin (**30**) [143] were reported. Arfan et al. have reported 11-O-galloylbergenin (**13**) possesses analgesic and antiinflammatory activities against formalin-induced noxious pains and carrageenan-induced paw oedema in rats, respectively [220]. It was also found more potent in the total antioxidant phosphomolybdate assay, reducing power assay and antiplasmodial assay [42]. Selective α -glucosidase inhibitory activity of compound **13** was reported [111]. 3,11-Di-O-galloylbergenin (**15**) has displayed a moderate antilipid

droplet accumulation activity [221]; 11-O-(4'-O-methylgalloyl)bergenin (**24**) inhibited arachidonic acid-induced platelet aggregation more efficiently than acetylsalicylic acid [222]; and bergecin B (**19**) showed potent inhibitory potential against the enzyme lipoxygenase ($IC_{50} = 24.3 \mu M$) [48]. Antiviral efficacy ($IC_{50} = 25 \mu g/mL$) of the dimer compound **35** was reported [27].

7. Derivatization of bergenin

7.1. Alkylation

Starting from bergenin (**1**), bergenin pentamethyl ether (**36**) was prepared using methyl iodide and freshly prepared Ag_2O in DMF [93]. 8,10-Di-O-methylbergenin (**7**) was synthesized by methylation of bergenin (**1**) and norbergenin (**2**) with diazomethane [97,139]. Shah et al. have prepared the alkyl derivatives of bergenin (**7**, **37-50**) by treating alkyl halides with bergenin (**1**) under mild basic conditions [107] (Fig. 3). The alkyl derivatives **7**, **40** and **50** showed nitric oxide inhibitory activity; **40** and **50** were $TNF-\alpha$ inhibitors; and **43**, **48** and **50** exhibited moderate antiinflammatory activity.




Fig. 3. Alkyl derivatives of bergenin.

7.2. Acylation

Several researchers have synthesized pentaacetylbergenin (**51**) by treating bergenin (**1**) with acetic anhydride or acetyl chloride in the presence of a base (such as pyridine, triethyl amine, DMAP, etc.) [5,49,50,74,93,99,124,162,225]. The acetylbergenin (**51**) has exerted hepatoprotective activity against D-galactosamine-induced cytotoxicity in cultured rat hepatocytes, and restored glutathione levels and decreased activities of glutathione S-transferase and glutathione reductase [226,99]. Antinociceptive activity of the acetylbergenin in mice was also reported [74]. Using different acid chlorides, bergenin acylates (**51-54**) were synthesized by Jung et al. [225] (Fig. 4). Comparing to the parent bergenin (**1**), these derivatives (**51-54**) possessed enhanced antiinflammatory activity (suppression of lipopolysaccharide (LPS)-induced nitric oxide generation) and antinarcotic effects on morphine dependence in mice.




Fig. 4. Preparation of bergenin pentaacylates [225].

Tiwari and Khosa have prepared bergenin diethyl ether triacetate (**55**) in two steps: ethylation of phenolic groups of bergenin followed by acetylation of alcoholic groups (Fig. 5) [162].




Fig. 5. Preparation of bergenin diethyl ether triacetate (**55**) [162].

Kumar et al. have synthesized bergenin monoacylates (**13**, **22**, **57**) by coupling of acid chlorides with dibenzyl bergenin (**56**) followed by hydrogenation (Fig. 6) [106]. Compounds **13** and **57** exhibited potent antiglycation activity with the IC₅₀ values of 12.28 and 60.75 μM, respectively. According to the authors, formation of fluorescent advanced glycation end products associated with secondary diabetes complications occurs in three stages – formation of Schiff-bases, Amadori rearrangement and decarbonyls formation together with cross linking with proteins. 11-O-Galloylbergenin (**13**) has been displayed potent inhibitory activities at all the three stages.




Fig. 6. Preparation of bergenin monoacylates [106].

7.3. Demethylation

Demethylation of bergenin (**1**) using HI yielded norbergenin (**2**) [139]. Norbergenin (**2**) was also synthesized by acetylation of bergenin followed by demethylation with BCl₃ and deprotection of hydroxyls [224,49] (Fig. 8). Pouységu et al. reported chemoselective oxygenative O-demethylation of phenolic methyl aryl ethers of bergenin (**1**) and its monobenzyl derivatives (**58**, **59**) (monobenzylation of the phenolic group of bergenin was achieved by treating with benzyl chloride in the presence of

NaHCO₃ and NaI in DMF) using stabilized 2-iodoxybenzoic acid (SIBX) afforded demethylated products norbergenin (**2**) and its monobenzylated catacholic derivatives (**60**, **61**) (Fig. 7) [116].




Fig. 7. Monobenylation of bergenin and demethylation [116].

7.4. Selective monoesterification

Takahashi et al. converted bergenin (**1**) into norbergenin (**2**) and then modified the sugar part by coupling with a variety of fatty acids (Fig. 8) [224]. Selective benzylation of the phenolic hydroxyl groups in norbergenin (**2**) afforded tribenzyl norbergenin (**62**). The hydroxyl groups on the sugar part were then esterified with various chain lengths of fatty acids to give compounds **63-68**. Monoesterification of the hydroxyl group at C-11 position was accomplished by coupling **62** with DCC and hexanoic acid followed by hydrogenation to obtain norbergenin 11-caproate (**69**). Norbergenin 4-caproate (**70**) was prepared by protection of hydroxyl groups at C-11 and C-3 as an acetal, esterification at free C-4 followed by hydrogenation. Norbergenin 3-caproate (**71**) was obtained by acid hydrolysis of compound **65**. These derivatives (**63-71**) greatly enhanced the free radical scavenging activity and prevented neuronal death of fetal rat cortical neurons, caused by reactive oxygen species, in Dulbecco's modified Eagle's medium (DMEM) supplemented with N2.




Fig. 8. Selective esterification [224].

Nazir et al. prepared bergenin pentaacetate (**51**) by acetylation of bergenin (**1**) and was subjected to lipase (PSL, lipase from *Pseudomonas cepacia* immobilized on Hyflo Super Gel or MML, lipase from *Mucor miehei* immobilized in Sol-Gel-AK)-catalyzed regioselective alcoholysis in dry n-butanol to obtain 3,4,10,11-tetraacetate of bergenin (**72**) (Fig. 9) [50]. The free-hydroxyl group of compound **72** was acylated with various fatty acids by reacting with DCC in the presence of DMAP to yield acyl derivatives **73-76**. Compounds **51** and **73** showed DPPH radical scavenging activity; compounds **51**, **72** and **73** showed considerable antibacterial activity; and compounds **75** and **76** were found to promote xanthine oxidase catalyzing ability against xanthine to produce uric acid.




Fig. 9. Acetylation, alcoholysis and esterification of bergenin [50].

Mozhaev et al. [227] demonstrated a combinatorial strategy for the sequential acylations of bergenin (**1**) employing immobilized lipases (Chirazyme L-2, Chirazyme L-9, lipase PS30 and lipase FAP-15) and protease (Subtilisin Carlsberg) at 11 and 4 positions, respectively (Fig. 10). Regioselective deacylation at 11 position was achieved when the diacylated product was hydrolysed with the lipase

in acetonitrile containing 2% water. A set of twelve acyl donors including vinyl and trifluoroethyl esters were used to prepare a library of 24 monoacylated and 144 diacylated derivatives of bergenin.




Fig. 10. Sequential enzymatic acylation/hydrolysis for production of acylated bergenin derivatives [227].

7.5. Esterification of the primary hydroxyl under Mitsunobu conditions

Kumar et al. synthesized bergenin esters (**77-84**) by coupling of aromatic acids with bergenin (**1**) employing Mitsunobu conditions and their potentiality for free radical ABTS^{++} scavenging were reported. (Fig. 11) [106].




Fig. 11. Synthesis of bergenin esters [106].

Bergenin esters (**21,22,26,27,77,86-89**) were prepared by Kashima and Miyazawa (Fig. 12), and their mushroom tyrosinase inhibitory and antioxidant activities were reported [40]. Among them compound 11-O-protocatechuoylbergenin (**87**) exhibited potent tyrosinase inhibitory activity ($\text{IC}_{50} = 17.5 \mu\text{M}$). These compounds were also evaluated for β -secretase (BACE1) activity [41]. Among them, compound **87** was the most potent inhibitor with an IC_{50} value of $0.6 \mu\text{M}$. Other analogy compounds **22**, **26** and **88** displayed IC_{50} values of $<10.0 \mu\text{M}$ in BACE1 inhibitory activity.




Fig. 12. Synthesis of bergenin esters [40].

In a subsequent paper, Kashima and Miyazawa have prepared di-O-methylbergenin analogues (**90-98**) starting from bergenin (**1**) and their antioxidant property and tyrosinase inhibitory activity were studied (Fig. 13) [228]. Among tested analogues, compound **98** exhibited a greater antioxidant activity and tyrosinase inhibitory activity ($\text{IC}_{50} = 9.1 \mu\text{M}$).




Fig. 13. Synthesis of di-O-methylbergenin esters [228].

7.6. Dimerization

Wang et al. have reported biotransformation of bergenin (**1**) with white rot fungus *Pleurotus ostreatus* to produce C–C coupled dimeric bergenin derivative **35** (structure shown in Fig. 2) [229]. The IC_{50}

value of the dimeric compound (2.13 mM/mL) was doubled than monomer bergenin (**1**) (1.07 mM/mL) in their *in vitro* antioxidant activity in DPPH assay.

7.7. Degradation

Bergenin (**1**) was degraded with *Erwinta herbicola*, a strain of soil bacteria isolated from the rhizosphere of *Bergenia crassifolia*, into 4-O-methylgallic acid [230]. Hattori et al. also reported degradation of bergenin (**1**) into 4-O-methylgallic acid by a mixture of human intestinal bacteria isolated from human feces [231].

8. Total synthesis

Some attentions in the synthesis of bergenin have been paid in the past due to its interesting biological activities. In 1958, Hay and Haynes have reported the first synthesis of bergenin (**1**) by employing the reaction of tetra-O-acetyl- α -D-glucopyranosyl bromide with methyl-4-O-methyl gallate in the presence of sodium methoxide [5]. Although it was a low yielding procedure, the authors could establish the structure of bergenin (**1**) through the laboratory synthesis. Thereafter for about four decades, not much work has been done on bergenin except some reports on its isolation from the plant sources. Some attempts to prepare bergenin (**1**) were not successful [232,233]. Bergenin has gained much attention from the beginning of 21st century, as its pharmacological properties were gradually explored.

In 1991, a ten step synthesis of 8,10-di-O-methylbergenin (**7**) has been reported by Schmidt and coworker in overall 5.2% yield starting from perbenzylated trifluoroacetyl glucose (**99**) (Fig. 14) [234]. In the presence of $\text{BF}_3 \cdot \text{OEt}_2$, the β -glucosyl trifluoroacetate (**99**) was coupled with 1,2,3-trimethoxybenzene affording 4- β -C-glucosylarene **100**. Removal of the benzyl groups from compound **100** by hydrogenolysis followed by O-methoxycarbonylation with methyl chloroformate produced compound **101**. Bromination of **101** gave the 1-bromo derivative **102**. Bromine/lithium exchange with butyl lithium generated the lithiated species **103** *in situ*, which was reacted with diphenyl disulphide to yield the phenyl sulphide **104**. Oxidation of **104** with 3-chloroperoxy benzoic acid gave the diastereomeric sulfoxide **105** in 1:2 ratio. After chromatographic separation, the major isomer was treated with lithium di-isopropylamide to generate the C-lithiated compound **106**, which was reacted with methyl chloroformate to give compound **107**. Raney nickel desulphurization then produced the prebergenin-type compound **108**, which after lactonization by treating with methanolic sodium methoxide produced 8,10-di-O-methylbergenin (**7**). Acetalization of alcohol functionality with acetic anhydride in pyridine afforded compound **109**.




Fig. 14. Synthesis of 8,10-di-O-methylbergenin (**7**) [234].

Synthesis of peracetate of 8,10-di-O-methylbergenin (**109**) was reported by Martin and coworker, based on an intramolecular C-glycosylation of a 2-(3',4',5'-trimethoxy)benzyl n-pentenyl glucoside followed by oxidation of the benzylic methylene group (Fig. 15) [235]. Starting from peracetylated glucosyl bromide **110**, the compound **109** was obtained in 8.5% yield over nine steps. Tetra-O-acetyl- α -D-glucopyranosyl bromide (**110**) under Lemieux-Morgan conditions provided O-pentenyl orthoester (**111**). The acetyl groups in compound **111** were replaced with benzyl groups and treated with trimethylsilyl triflate followed by a base treatment afforded pentenyl β -glucoside **112**. O-Benzoylation of **112** using 3,4,5-trimethoxybenzyl chloride provided compound **113**. The treatment of **113** with iodonium dicollidine perchlorate (IDCP) promoted intramolecular C-arylation affording **114**. The treatment of **114** with BF₃·Et₂O resulted epimerization. The resulting product **115** was deprotected by hydrogenolysis and reacetylated. The benzylic position of **116** was oxidized using catalytic amount of ruthenium tetroxide to give the final lactone **109**.




Fig. 15. Synthesis of peracetate of 8,10-di-O-methylbergenin (**109**) [235].

A short, five step synthesis of 8,10-di-O-methylbergenin (**7**) was reported by Seeberger and coworkers (Fig. 16) [236]. In the presence of trimethylsilyl trifluoromethane sulphonate (TMSOTf), 2,3,4,6-tetra-O-benzyl glucopyranosyl trichloroacetimidate (**117**) and perbenzylated glucosyl diphenyl phosphate **118** reacted with 3,4,5-trimethoxy phenol to produce the β -configured C-glycoside **119**. Triflate **120**, which was obtained upon treatment with triflic anhydride/lutidine, was subjected to Pd(0)-catalyzed aryl carbonation affording C-glucosyl benzoic acid derivative **121**. Hydrogenation of **121** with Pearlman's catalyst afforded debenzylated product. Finally, 8,10-di-O-methylbergenin (**7**) was obtained upon treatment of the debenzylated product with SOCl₂ in methanol.




Fig. 16. Synthesis of 8,10-di-O-methylbergenin (**7**) [236].

Sakamaki and coworkers have synthesized 8,10-di-O-methylbergenin (**7**) by application of the aryl- β -C-glycosidation method (Fig. 17) [237]. Aryl bromide **123**, obtained by bromination of commercial methyl 3,4,5-trimethoxybenzoate (**122**), underwent Pd-catalyzed coupling reaction with glucal boronate (**124**). After hydroboration-oxidation, aryl- β -C-glucoside **126** was obtained. Compound **126** was oxidized with MnO₂ to produce lactone **127**. Deprotection of the silyl function afforded lactone-

ring opening product, which was cyclized in the presence of thionyl chloride to yield 8,10-di-O-methylbergenin (**7**).




Fig. 17. Synthesis of 8,10-di-O-methylbergenin (**7**) [237].

While preparing this manuscript, the first total synthesis of bergenin (**1**) was reported by Parkan and coworkers (Fig. 18) [238]. Hydroboration-oxidation of the Suzuki-Miyaura cross-coupling product **129** furnished arylglucoside **130** diastereoselectively. Selective oxidation of the benzyl alcohol moiety was accompanied by cyclization to produce lactone **132**. Removal of silyl protecting group followed by reductive debenzoylation afforded bergenin (**1**). Thus bergenin was synthesized in six steps from bromide **128** in 40% overall yield.




Fig. 18. Synthesis of bergenin (**1**) [238].

9. Conclusion

About 25% of medicines prescribed worldwide today are derived from plants and about 80% of the people living in the developing countries still rely on traditional plant-derived medicines [239]. Among the plant-derived medicines, bergenin (**1**) is considered as an antitussive agent. Literature

review reveals that bergenin exhibits antiviral, antifungal, antitussive, antiplasmodial, antiinflammatory, antihepatotoxic, antiarrhythmic, antitumor, antiulcerogenic, antidiabetic and wound healing properties. It displayed poor antimicrobial and antioxidant activities. In Ayurvedic formulations, bergenin containing plants (particularly *Bergenia ligulata*) are used to dissolve urinary calculi and the traditional use of the plant materials was also supported by the experimental results [240,241]; however, bergenin itself exhibited less significant antiurolithiatic activity [193]. Therefore, efficient antiurolithiatic constituents present in genus *Bergenia* are essential to be searched. It can be concluded that bergenin is an important secondary metabolite that responsible for multiple actions for the betterment of human health. Bergenin is abundantly distributed in genera Saxifragaceae, Euphorbiaceae, Myrsinaceae, Dipterocarpaceae and Fabaceae. Up to 9.77% of bergenin was estimated to be deposited in the rhizome of *Bergenia purpurascens* (Saxifragaceae) [200]. Several bergenin derivatives were isolated from plants, synthesized in laboratories and studied their biological activities. Among them, norbergenin (**2**) is a most important and potentially bioactive constituent. Although bergenin was first isolated from the rhizomes of *Saxifraga (Bergenia) siberica* in 1881 [11], its structure was confirmed only in 1958 [5-7]. Bergenin has gained importance in scientific community only in 21st century when its pharmacological properties were gradually explored. As a consequence, a number of pharmacokinetic studies, analyses, isolation techniques and total synthesis of bergenin and its derivatives were recently reported. In conclusion, this article provides a comprehensive review on bergenin and its derivatives comprising all the available literature from the beginning and may help in development of new drugs in the future.

Conflicts of interest

The author has no conflicts of interest to declare.

Acknowledgements

Nepal Academy of Science and Technology (NAST) is acknowledged for essential support. I thank Dr. Yugesh Kharel, Dr. Sajan L. Shyaula and Dr. Achyut Adhikari for literature collection.

References

- [1] Taneyama M, Yoshida S. Studies on C-glycosides in higher plants II. Incorporation of ¹⁴C-glucose into bergenin and arbutin in *Saxifraga stolonifera*. Bot Mag Tokyo 1979;92:69–73.
- [2] Lu X, Wang J. Advances in the study of *Bergenia* plants. Zhong Yao Cai (= Journal of Chinese Medicinal Materials) 2003;26:58–60.
- [3] Tschitschibabin AE, Kirssanow AW, Korolew AJ, Woroschzow jun NN. Über nichtgerbende substanzen des extraktes aus dem wurzelstock des badans (*Saxifraga crassifolia*). I. Bergenin. Justus Liebigs Annalen der Chemie 1929;469:93–127.
- [4] Shimokôriyama M. Science (Japan) 1950;20:576–7 (cited in Ref. 5).
- [5] Hay JE, Haynes LJ. Bergenin, a C-glycopyranosyl derivative of 4-O-methylgallic acid. J Chem Soc 1958;2231–8.
- [6] Posternak, Th, Dürr K. Sur la constitution de la bergénine. Helvetica Chimica Acta 1958;41:1159–1162.
- [7] Fujise S-I, Suzuki M, Watanabe Y, Matsueda S. Studies of the structure of bergenin. Bull Chem Soc Jpn 1959;32:97–8.
- [8] Frick W, Hofmann J, Fischer H, Schimdt RR. The structure of bergenin. Carbohydrate Res 1991;210:71–7.
- [9] Caldas CS, De Simone CA, Pereira MA, Malta VRS, Carvalho RLP, Da Silva TBC et al. Bergenin monohydrate, a constituent of *Humiria balsamifera*, at 120 K. Acta Crystallographica Sec E 2002;58:609–611.
- [10] Ye Y-P, Sun H-X, Pan Y-J. Bergenin monohydrate from the rhizome of *Astilbe chinensis*. Acta Crystallographica Sec C 2004;60:397–8.
- [11] De Morelle ME, Chatin M. Sur un nouvel hydrate de carbone. Compt Rend 1881;93:646.

- [12] Olennikov DN, Chekhirova GV. 6"-Galloylpicein and other phenolic compounds from *Arctostaphylos uva-ursi*. Chem Nat Prod 2013;49:1–7.
- [13] Sumino M, Sekine T, Ruangrunsi N, Igarashi K, Ikegami F. Ardisiphenols and other antioxidant principles from the fruits of *Ardisia colorata*. Chem Pharm Bull 2002;50:1484–7.
- [14] Ni M, Han L. Studies on the chemical constituents of *Ardisia crenata* Sims. Bull Chinese Materia Medica 1988;59:33–4.
- [15] Jia Z, Mitsunaga K, Koike K, Ohmoto T. New bergenin derivatives from *Ardisia crenata*. Natural Medicines 1995;49:187–9.
- [16] Deng J, Xiao X, Tong X, Li G. Preparation of bergenin from *Ardisia crenata* Sims and *Rodgersia sambucifolia* Hemsl based on microwave-assisted extraction/high-speed counter-current chromatography. Separation and Purification Technology 2010;74:155–9.
- [17] Liu N, Li Y, Gua J-X, Qian D-G. Studies on the taxonamy of the genus *Ardisia* (Myrsinaceae) from China and the occurrence and quantity of bergenin in the genus. Acta Academiae Medicinae Shanghai 1993;20:49–54.
- [18] Mu L-H, Feng J-Q, Liu P. A new bergenin derivative from the rhizome of *Ardisia gigantifolia*. Nat Prod Res 2013;27:1242–5.
- [19] Piacente S, Pizza C, Tommasi ND, Mahmood N. Constituents of *Ardisia japonica* and their in vitro anti HIV activity. J Nat Prod 1996;59:565–9.
- [20] Li Y-F, Hu L-H, Lou F-C, Li J, Shen Q. PTP1B inhibitors from *Ardisia japonica*. J Asian Nat Prod Res 2005;17:13–8.
- [21] Ma C-F, Luo M, Lin L-M, Li C, Wang Z-M, Cheng Y-Y. Chemical constituents of *Ardisia punctata*. Zhongguo Zhongyao Zazhi (= China Journal of Chinese Materia Medica) 2012;37:3422–5.
- [22] Su Y, Xu J-J, Bi J-L, Wang Y-H, Hu G-W, Yang J et al. Chemical constituents of *Arisaema franchetianum* tubers. J Asian Nat Prod Res 2013;15:71–7.
- [23] Sun HX, Ye YP, Yang K. Studies on the chemical constituents in radix *Astilbe chinensis*. Chinese J Chinese Materia Medica 2003;27:751–4.
- [24] Xue Y, Xu X-M, Yan J-F, Deng W-L, Liao X. Chemical constituents from *Astilbe chinensis*. J Asian Nat Prod Res 2011;13:188–191.
- [25] Zou Y, Cui Y. Chemical constituents from *Astilbe myriantha*. Zhongyaocai (= Journal of Chinese Medicinal Materials) 2012;35:1095–7.
- [26] Tandon M, Shukla YN, Tripathi AK, Sharma S. Antifeedant activity of bergenin isolated from *Astilbe rivularis*. Fitoterapia 1996;67:277–8.
- [27] Rajbhandari M, Lalk M, Mentel R, Lindequist U. Antiviral activity and constituents of the Nepalese medicinal plant *Astilbe rivularis*. Rec Nat Prod 2011;5:138–142.
- [28] Han L-K, Ninomiya H, Taniguchi M, Baba K, Kimura Y, Okuda H. Norepinephrine-augmenting lipolytic effectors from *Astilbe thunbergii*. J Nat Prod 1998;61:1006–1011.
- [29] Kimura Y, Sumiyoshi M, Sakanaka M. Effects of *Astilbe thunbergii* rhizomes on wound healing: Part 1. Isolation of promotional effectors from *Astilbe thunbergii* rhizomes on burn wound healing. J Ethnopharmacol 2007;109:72–7.
- [30] Taneyama M, Yoshida S. Studies on C-glycosides in higher plants I. Occurrence of bergenin in Saxifragaceae. J Plant Res 1978;91:109–112.
- [31] Ostrowska B, Gorecki P. Investigations on possibility of utilization of *Bergenia* leaves to therapeutics in place of arbutin and tannin raw materials deficiency. 1. Phytochemical investigations [*Bergenia crassifolia* and *Bergenia cordifolia*]. Herba Polonica 1988;34:21–5.
- [32] Ostrowska B, Gorecki P, Wolska D. Investigation on possibility of utilization of *Bergenia* leaves to therapeutics in place of arbutin and tannin raw materials deficiency. Part 2. Isolation of bergenin and a method of its quantitative determination [*Bergenia crassifolia*, *Bergenia cordifolia*]. Herba Polonica 1989;35:117–122.
- [33] Roselli M, Lentini G, Habtemariam S. Phytochemical, antioxidant and anti- α -glucosidase activity evaluations of *Bergenia cordifolia*. Phytother Res 2012;26:908–914.
- [34] Sadikov VS, Guthner RA. Zur Kenntnis des bergenins. I. Bergenin. Biochemische Zeitschrift 1927;190:340–351.

- [35] Pozharitskaya ON, Ivanova SA, Shikov AN, Makarov VG, Galambosi B. Separation and evaluation of free radical-scavenging activity of phenol components of green, brown, and black leaves of *Bergenia crassifolia* by using HPTLC-DPPH• Method. J Sep Sci 2007;30:2447–2451.
- [36] Bahl CP, Murari R, Parthasarathy MR, Seshadri TR. Components of *Bergenia strecheyi* and *B. ligulata*. Indian J Chem 1974;12:1038–9.
- [37] Jain MK, Gupta K. Isolation of bergenin from *Saxifraga ligulata* Wall. J Indian J Chem Soc 1962;39:559–560.
- [38] Kohlmunzer S, Chojnacka-Wojcik E. Isolation of bergenin and some of its pharmacological properties. Bull Liaison Groupe Polyphenols 1986;13:590–3.
- [39] Reddy UDC, Chawla AS, Deepak M, Singh D, Handa SS. High performance liquid chromatographic determination of bergenin and (+)-afzelechin from different parts of Paashaanbhed (*Bergenia ligulata* Yeo). Phytochem Anal 1999;10:44–7.
- [40] Kashima Y, Miyazawa M. Synthesis and biological evaluation of bergenin analogues as mushroom tyrosinase inhibitors. Arch Pharm Res 2012;35:1533–1541.
- [41] Kashima Y, Miyazawa M. Structure-activity relationships for bergenin analogues as β -secretase (BACE1) inhibitors. J Oleo Sci 2013;62:391–401.
- [42] Uddin G, Sadat A, Siddiqui BS. Comparative antioxidant and antiplasmodial activities of 11-O-galloylbergenin and bergenin isolated from *Bergenia ligulata*. World Applied Sciences Journal 2013;27:977–981.
- [43] Xin-Min C, Yoshida T, Hatano T, Fukushima M, Okuda T. Galloylarbutin and other polyphenols from *Bergenia purpurascens*. Phytochemistry 1987;26:515–7.
- [44] Bajracharya GB, Maharjan R. Development of an ease methodology for the isolation of Ayurveda drug bergenin. J Nepal Chem Soc 2013;32(1):65–69.
- [45] Cui Y. Chemical constituents from rhizomes of *Bergenia scopulosa* (II). Zhongcaoyao (= Chinese Traditional and Herbal Drugs) 2012;43:1704–7.
- [46] Wei Y-Y. Chemical constituents from *Bergenia scopulosa* (I). Zhongguo Shiyang Fangjixue Zazhi (= Chinese Journal of Experimental Medical Formulae) 2012;18:154–6.
- [47] Chowdhary S, Kumar H, Verma DL. Chemical examination of *Bergenia stracheyi* (Hk.) for antioxidative flavonoids. Nature Sci 2009;7:29–34.
- [48] Siddiq F, Fatima I, Malik A, Afza N, Iqbal L, Lateef M et al. Biologically active bergenin derivatives from *Bergenia stracheyi*. Chemistry and Biodiversity 2012;9:91–8.
- [49] Nazir N, Koul S, Qurishi MA, Taneja SC, Ahmad SF, Bani S et al. Immunomodulatory effect of bergenin and norbergenin against adjuvant-induced arthritis – A flow cytometric study. J Ethnopharmacol 2007;112:401–5.
- [50] Nazir N, Koul S, Qurishi MA, Najar MH, Zargar MI. Evaluation of antioxidant and antimicrobial activities of bergenin and its derivatives obtained by chemoenzymatic synthesis. European J Med Chem 2011;46:2415–2420.
- [51] Zhao J, Zeng L-H, Li X, Dong X-P, Yan Y-M, Cheng Y-X. Brachystemols A-C, three new furan derivatives from *Brachystemma calycinum*. J Asian Nat Prod Res 2011;13:915–9.
- [52] Wei X-H, Yang S-J, Liang N, Hu D-Y, Jin L-H, Xue W et al. Chemical constituents of *Caesalpinia decapetala* (Roth) Alston. Molecules 2013;18:1325–1336.
- [53] Chaudhary GR, Sharma VN, Dar ML. Chemical examination of the roots of *Caesalpinia digyna*. J Sci Ind Res 1954;13B:147–8.
- [54] Srinivasan R, Chandrasekar MJN, Nanjan MJ, Suresh B. Antioxidant activity of *Caesalpinia digyna* root, J Ethnopharmacol 2007;113:284–291.
- [55] Singh U, Kunwar A, Srinivasan R, Nanjan MJ, Priyadarsini KI. Differential free radical scavenging activity and radioprotection of *Caesalpinia digyna* extracts and its active constituents. J Radiat Res 2009;50:425–433.
- [56] Kumar R, Patel DK, Prasad SK, Laloo D, Krishnamurthy S, Hemalatha S. Type 2 antidiabetic activity of bergenin from the roots of *Caesalpinia digyna* Rottler. Fitoterapia 2012;83:395–401.
- [57] Roy SK, Agrahari UC, Gautam R, Srivatava A, Jachak SM. Isointracatinol, a new antioxidant homoisoflavonoid from the roots of *Caesalpinia digyna* Rottler. Nat Prod Res 2012;26:690–5.

- [58] Yodsaoue O, Karalai C, Ponglimanont C, Tewtrakul S, Chantrapromma S. Potential anti-inflammatory diterpenoids from the roots of *Caesalpinia mimosoides* Lamk. *Phytochemistry* 2010;71:1756–1764.
- [59] De Oliveira CM, Nonato FR, De Lima FO, Couto RD, David JP, David JM et al. Antinociceptive properties of bergenin. *J Nat Prod* 2011;74:2062–8.
- [60] Alves CQ, David JM, David JP, Villareal CF, Soares MBP, De Queiroz LP et al. Flavonoids and other bioactive phenolics isolated from *Cenostigma macrophyllum* (Leguminosae). *Química Nova* 2012;35:1137–1140
(http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0100-40422012000600013).
- [61] Cao P, Pu X-F, Peng S-L, Zhang X-R, Ding L-S. Chemical constituents from *Cimicifuge foetida*. *J Asian Nat Prod Res* 2005;7:145–9.
- [62] Li D-S, Nian Y, Sun Y, Qiu M-H. Three new cycloartane (= 9,19-cycloanostane) glycosides from *Cimicifuge foetida*. *Helvetica Chimica Acta* 2011;94:632–8.
- [63] Aiyar SN, Jain MK, Krishnamurti M, Seshadri TR. Chemical components of the roots of *Connarus monocarpus*. *Phytochemistry* 1964;3:335–9.
- [64] Kim MH, Ha SY, Oh MH, Kim HH, Kim SR, Lee MW. Anti-oxidative and anti-proliferative activity on human prostate cancer cells lines of the phenolic compounds from *Corylopsis coreana* Uyeki. *Molecules* 2013;18:4876–4886.
- [65] Hattori S. Corylopsin, a crystalline constituent of the bark of *Corylopsis spicata*. *Acta Phytochim (Japan)* 1929;4:327–341.
- [66] Li C, Chen X, Fang D, Li G. A new bergenin derivative from *Corylopsis willmottiae*. *Chemistry of Natural Compounds* 2011;47:194–6.
- [67] Musgrave OC, Skoyles D. Ebenaceae extractives. Part IV. Diosindigo A, a pigment from several *Diospyros* species. *J Chem Soc Perkin Trans 1* 1974;1128–1131.
- [68] Ito T, Tanaka T, Iinuma M, Nakaya K-I, Takahashi Y, Sawa R et al. Two new resveratrol (= 5-[(1E)-2-(4-hydroxyphenyl)ethenyl]-benzene-1,3-diol) tetramers with a tetrahydrofuran ring from *Dipterocarpus grandiflorus*. *Helvetica Chimica Acta* 2004;87:479–495.
- [69] Wibowo A, Ahmat N, Hamzah AS, Sufian AS, Ismail NH, Ahmad R et al. Malaysianol A, a new trimer resveratrol oligomer from the stem bark of *Dryobalanops aromatic*. *Fitoterapia* 2011;82:676–681.
- [70] Wibowo A, Ahmat N, Hamzah AS, Ismail NH, Ahmad R, Jaafar FM. Resveratrol oligomers from the stem bark of *Dryobalanops aromatic*. *Biochemical Systematics and Ecology* 2012;40:62–4.
- [71] Kuspradini H, Mitsunaga T, Mihara R, Ohashi H. Investigating glucosyltransferase inhibitory activities of polyphenols from kapur (*Dryobalanops* sp.) heartwood extracts. *J Nat Med* 2007;61:462–7.
- [72] Da Silva SL, De Oliveira VG, Yano T, Nunomura R de CS. Antimicrobial activity of bergenin from *Endopleura uchi* (Huber) Cuatrec. *Acta Amazonica* 2009;39:187–192.
- [73] Nunomura RCS, Oliveira VG, Da Silva SL, Nunomura SM. Characterization of bergenin in *Endopleura uchi* bark and its anti-inflammatory activity. *J Braz Chem Soc* 2009;20:1060–4.
- [74] Borges JCM, Ripardo Filho H da S, Guilhon GMSP, Carvalho JCT, Santos LS, Sousa PJC. Antinociceptive activity of acetylbergenin in mice. *Lat Am J Pharm* 2011;30:1303–8.
- [75] Magalhães LA, Lima MP, Marinho HA, Ferreira AG. Identificação de bergenina e carotenóides no fruto de uchi (*Endopleura uchi*, Humiriaceae). *Acta Amazonica* 2007;37:447–450.
- [76] Li RW, Leach DN, Myers SP, Lin GD, Leach GJ, Waterman PG. A new anti-inflammatory glucoside from *Ficus racemosa* L. *Planta Medica* 2004;70:421–6.
- [77] Lin GD, Li RW, Myers SP, Leach DN. A method of selecting plants with anti-inflammatory potential for pharmacological study. *Nat Prod Commun* 2008;3:71–6.
- [78] Ahmed F, Siddesha JM, Urooj A, Vishwanath BS. Radical scavenging and angiotensin converting enzyme inhibitory activities of standardized extracts of *Ficus racemosa* stem bark. *Phytother Res* 2010;24:1839–1843.
- [79] Jain R, Rawat S, Jain SC. Phytochemicals and antioxidant evaluation of *Ficus racemosa* root bark. *J Pharm Res (Guragaon India)* 2013;6:615–9.
- [80] Ahmad SA, Kapoor SK, Zaman A. Bergenin in *Flueggea microcarpa*. *Phytochemistry* 1972;11:452.
- [81] Kumar S, Sahai M, Ray AB. Chemical constituents of the leaves of *Flueggea microcarpa*. *Planta Medica* 1985;59:466.

- [82] Jahromi MAF, Chansouria JPN, Ray AB. Hypolipidaemic activity in rats of bergenin, the major constituent of *Flueggea microcarpa*. *Phytother Res* 1992;6:180–3.
- [83] Pu H-L, Huang X, Zhao J-H, Hong A. Bergenin is the antiarrhythmic principle of *Fluggea virosa*. *Planta Medica* 2002;68:372–4.
- [84] Nyasse B, Nono J, Sonke B, Denier C, Fontaine C. Trypanocidal activity of bergenin, the major constituent of *Flueggea virosa*, on *Trypanosoma brucei*. *Pharmazie* 2004;59:492–4.
- [85] Wang G-C, Liang J-P, Wang Y, Li Q, Ye W-C. Chemical constituents from *Flueggea virosa*. *Chinese Journal of Natural Medicines* 2008;6:251–3.
- [86] Alkadi KAA, Adam A, Taha M, Hasan MH, Shah SAA. Antiplatelet aggregation activity of 5-hydroxyfalcone, 2'-hydroxyflavanone, paeonol and bergenin isolated from stem bark of *Garcinia malaccensis* in human whole blood. *Oriental Journal of Chemistry* 2013;29 (<http://www.orientjchem.org/?p=262>).
- [87] Lu S, Zhang G. Alkaloids from *Gendarussa vulgaris* Nees. *Nat Prod Res* 2008;22:1610–3.
- [88] Thang TD, Kuo P-C, Yu C-S, Shen Y-C, Hoa LTM, Thanh TV et al. Chemical constituents of the leaves of *Glochidion obliquum* and their bioactivity. *Arch Pharm Res* 2011;34:383–9.
- [89] Takeda Y, Mima C, Masuda T, Hirata E, Takushi A, Otsuka H. Glochidioboside, a glucoside of (7S,8R)-dihydrodehydrodiconiferyl alcohol from leaves of *Glochidion obovatum*. *Phytochemistry* 1998;49:2137–9.
- [90] Nasser JA, Yaacob WA, Din LB, Yamin BM, Latip J. Isolation of atranorin, bergenin and goniothalamin from *Hopea sangal*. *ARNP Journal of Engineering and Applied Sciences* 2009;4:92–5 (www.arnpjournals.com/jeas/research_papers/rp.../jeas_0209_163.pdf).
- [91] Tanaka T, Ito T, Ido Y, Nakaya K-I, Iinuma M, Chelladurai V. Hopeafuran and a C-glucosyl resveratrol isolated from stem wood of *Hopea utilis*. *Chem Pharm Bull* 2001;49:785–7.
- [92] Da Silva TBC, Alves VL, Mendonça LVH, Conserva LM, Da Rocha EMM, Andrade EHA et al. Chemical constituents and preliminary antimalarial activity of *Humiria balsamifera*. *Pharmaceutical Biol* 2004;42:94–7.
- [93] Ramaiah PA, Row LR, Reddy DS, Anjaneyulu ASR, Ward RS, Pelter A. Isolation and characterization of bergenin derivatives from *Macaranga peltata*. *J Chem Soc Perkin Trans 1* 1979;2313–6.
- [94] Minh CV, Thanh NTK, Quang TH, Cuong NX, Thin NN, Nam NH et al. Two new megastigmane sulphonoglucosides from *Mallotus anisopodus*. *Nat Prod Commun* 2009;4:889–892.
- [95] Shibata K, Shimogori M. *Nippon Kagaku Zasshi* 1949;70:36 (cited in: Noda T, Take T, Watanabe T, Abe J. The characterization of malloprenol and its ester with linolenic acid isolated from the leaves of *Mallotus japonicus*. *Bull Chem Soc Jpn* 1970;43:2174–6.)
- [96] Hasegawa T. Constituents of *Mallotus japonicus*. I. II. III. *Yakugaku Zasshi* (= Journal of the Pharmaceutical Society of Japan) 1941;61:307–320.
- [97] Yoshida T, Seno K, Takama Y, Okuda T. Bergenin derivatives from *Mallotus japonicus*. *Phytochemistry* 1982;21:1180–2.
- [98] Saijo R, Nonaka G-I, Nishioka I. Gallic acid esters of bergenin and norbergenin from *Mallotus japonicus*. *Phytochemistry* 1990;29:267–270.
- [99] Lim H-W, Kim H-S, Kim S-H, Chang M-J, Rhee GS, Choi J. Protective effects of acetylbergenin against carbon tetrachloride-induced hepatotoxicity in rats. *Arch Pharm Res* 2001;24:114–8.
- [100] Lim HK, Kim HS, Choi HS, Choi JW. Protective and therapeutic effects of *Malloti cortex* extract on carbon tetrachloride- and galactosamine-induced hepatotoxicity in rats. *J Appl Pharmacol* 1999;7:35–43.
- [101] Kim H-S, Lim H-K, Chung M-W, Kim YC. Antihepatotoxic activity of bergenin, the major constituent of *Mallotus japonicus*, on carbon tetrachloride-intoxicated hepatocytes. *J Ethnopharmacol* 2000;69:79–83.
- [102] Zhang Y-H, Fang L-H, Lee M-K, Ku B-S. In vitro inhibitory effects of bergenin and norbergenin on bovine adrenal tyrosine hydroxylase. *Phytother Res* 2003;17:967–9.
- [103] Mai NT, Cuong NX, Thao NP, Nam NH, Khoi NH, Minh CV et al. A new lignan dimer from *Mallotus philippensis*. *Nat Prod Commun* 2010;5:423–6.
- [104] Bandopadhyay M, Dhingra VK, Mukerjee SK, Pardeshi NP, Seshadri TR. Triterpenoid and other components of *Mallotus philippinensis*. *Phytochemistry* 1972;11:1511.

- [105] Arfan M, Amin H, Khan I, Shah MR, Shah H, Khan AZ et al. Molecular simulations of bergenin as a new urease inhibitor. *Med Chem Res* 2011;21:2454–7.
- [106] Kumar TV, Tiwari AK, Robinson A, Babu KS, Kumar RSC, Kumar DA et al. Synthesis and antiglycation potentials of bergenin derivatives. *Bioorg Med Chem Lett* 2011;21:4928–4931.
- [107] Shah MR, Arfan M, Amin H, Hussain Z, Qadir MI, Choudhary MI et al. Synthesis of new bergenin derivatives as potent inhibitors of inflammatory mediators NO and TNF- α . *Bioorg Med Chem Lett* 2012;22:2744–7.
- [108] Huang P-L, Wang L-W, Lin C-N. New triterpenoids of *Mallotus repandus*. *J Nat Prod* 1999;62:891–2.
- [109] Tomizawa S, Asume K, Suguro N. Bergenin: Isocoumarin from the stems of *Mallotus repandus*. *Phytochemistry* 1976;15:328.
- [110] Rana VS, Rawat MSM, Pant G, Nagatsu A. Chemical constituents and antioxidant activity of *Mallotus roxburghianus* leaves. *Chemistry and Biodiversity* 2005;2:792–8.
- [111] Habtemariam S, Cowley RA. Antioxidant and anti- α -glucosidase compounds from the rhizome of *Peltiphyllum peltatum* (Torr.) Engl. *Phytother Res* 2012;26:1656–1660.
- [112] Izawa K, Nagai M, Inoue T. Constituents of saxifragaceous plants. III. Triterpene acids and bergenin in *Peltoboykinia watanabei* and *Boykinia lycoctonifolia*. *Phytochemistry* 1973;12:1508.
- [113] Mebe PP, Makuhunga P. 11-(E)-p-Coumaric acid ester of bergenin from *Peltophorum africanum*. *Phytochemistry* 1992;31:3286–7.
- [114] Bessong PO, Obi CL, Andréola M-L, Rojas LB, Pouységu L, Igumbor E et al. Evaluation of selected South African medicinal plants for inhibitory properties against human immunodeficiency virus type 1 reverse transcriptase and integrase. *J Ethnopharmacol* 2005;99:83–91.
- [115] Theo A, Masebe T, Suzuki Y, Kikuchi H, Wada S, Obi CL et al. *Peltophorum africanum*, a traditional South African medicinal plant, contains an anti HIV-1 constituent, betulinic acid. *Tohoku J Exp Med* 2009;217:93–9.
- [116] Pouységu L, Sylla T, Garnier T, Rojas LB, Charris J, Deffieux D et al. Hypervalent iodine-mediated oxygenative phenol dearomatization reactions. *Tetrahedron* 2010;66:5908–5917.
- [117] Sulochana V, Sastry KNS, Rao VSS, Reddy KK. Isolation of bergenin from *Peltophorum ferruginum*. *Leather Science (Madras)* 1970;17:327.
- [118] Joshi BS, Kamat VN. Identity of peltophorin with bergenin. *Naturwissenschaften* 1969;56:89–90.
- [119] Swarnalakshmi T, Sethuraman MG, Sulochana N, Arivudainambi R. A note on the anti-inflammatory activity of bergenin. *Current Science* 1984;53:917 (www.ias.ac.in/jarch/currsci/53/00000992.pdf).
- [120] Raj MK, Duraipandian V, Agastian P, Ignacimuthu S. Antimicrobial activity of bergenin isolated from *Peltophorum pterocarpum* DC. flowers. *Asian Pacific Journal of Tropical Biomedicine* 2012;S901–4.
- [121] Zhang J, Nishimoto Y, Tokuda H, Suzuki N, Yasukawa K, Kitdamrongtham W et al. Cancer chemopreventive effect of bergenin from *Peltophorum pterocarpum* wood. *Chemistry and Biodiversity* 2013;10:1866–1875.
- [122] Folefoc GN, Bisseck JP, Fomum ZT, Bodo B. Constituents from the roots of *Pentaclethra macrophylla*. *Biochemical Systematics and Ecology* 2005;33:1280–2.
- [123] Jamal AK, Yaacob WA, Din LB. A chemical study on *Phyllanthus columnaris*. *European J Scientific Res* 2009;28:76–81.
- [124] Tanaka R, Matsunaga S. Triterpene dienols and other constituents from the bark of *Phyllanthus flexuosus*. *Phytochemistry* 1988;27:2273–7.
- [125] Priya OS, Viswanathan MBG, Balakrishna K, Venkatesan M. Chemical constituents and in vitro antioxidant activity of *Phyllanthus wightianus*. *Nat Prod Res* 2011;25:949–958.
- [126] Venkateswarlu K, Satyalakshmi G, Suneel K, Reddy TS, Raju TV, Das B. A benzofuranoid and two clerodane diterpenoids from *Pulicaria wightiana*. *Helvetica Chimica Acta* 2008;91:2081–8.
- [127] Cuong TD, Hung TM, Kim J-C, Huh J-I, Kwack SJ, Kang TS et al. Two new lignans from the roots of *Pulsatilla koreana*. *Planta Medica* 2011;77:66–9.
- [128] Zamarrud, Ali I, Hussain H, Ahmad VU, Qaiser M, Aryn A et al. Two new antioxidant bergenin derivatives from the stem of *Rivea hypocrateriformis*. *Fitoterapia* 2011;82:722–5.
- [129] Shen XW, Zhen SZ, Fu ZS, Chen Y. Isolation and identification of the chemical constituents of Chinese medicinal herb *Rodgersia aesculifolia* Batal. *Chemical Journal of Chinese Universities* 1987;6:528–532.

- [130] Wang Y, Jin Y, Liu Z. Optimization of the extraction process of bergenin from *Rodgersia pinnata*. Dali Xueyuan Xuebao (= Journal of Dali University) 2012;11:8–10.
- [131] Ogan AU. An isocoumarin from the bark of *Sacoglottis gabonensis*. Phytochemistry 1971;10:2832–3.
- [132] Faparusi SI, Bassir O. Effects of the extracts of the bark of *Sacoglottis gabonensis* on the microflora of palmwine. Appl Microbiol 1972;24:853–6.
- [133] Ekong DE, Ejike C. Antioxidant principles from the bark of *Spondianthus preussi* and *Sacoglottis gabonensis* (Family Humiriaceae). Journal of West African Science Association 1974;19:63–8.
- [134] Maduka HCC, Okoye ZSC, Eje A. The influence of *Sacoglottis gabonensis* stem bark extract and its isolate bergenin, Nigerian alcoholic beverage additives, on the metabolic and haematological side effects of 2,4-dinitrophenyl hydrazine-induced tissue damage. Vascular Pharmacology 2003;39:317–324.
- [135] Maduka HCC, Okoye ZSC. A Nigerian alcoholic beverage additive from *Sacoglottis gabonensis* as an antioxidant protector of mammalian cells against 2,4-dinitrophenyl hydrazine-induced lipid peroxidation. The Internet Journal of Toxicology 2006;3 (doi: 10.5580/15da).
- [136] De Abreu HA, Lago IA dos S, Souza GP, Piló-Veloso D, Duarte HA, Alcântara AF de C. Antioxidant activity of (+)-bergenin – A phytoconstituent isolated from the bark of *Sacoglottis uchi* Huber (Humiriaceae). Org Biomol Chem 2008;6:2713–8.
- [137] Zuo G-Y, Li Z-Q, Chen L-R, Xu X-J. In vitro anti-HCV activities of *Saxifraga melanocentra* and its related polyphenolic compounds. Antiviral Chemistry and Chemotherapy 2005;16:393–8.
- [138] Morita N, Shimizu M, Arisawa M, Koshi M. Studies on the medicinal resources. XXXVI. The constituents of the leaves of *Saxifraga stolonifera* Meerburg (Saxifragaceae). Chem Pharm Bull 1974;22:1487–9.
- [139] Taneyama M, Yoshida S, Kobayashi M, Hasegawa M. Isolation of norbergenin from *Saxifraga stolonifera*. Phytochemistry 1983;22:1053–4.
- [140] Chen Z, Liu Y-M, Yang S, Song B-A, Xu G-F, Bhadury PS et al. Studies on the chemical constituents and anticancer activity of *Saxifraga stolonifera* (L) Meeb. Bioorg Med Chem 2008;16:1337–1344.
- [141] Xian C, Gong X, Zhao C, Zhou X, Yang Z, Wang L. Chemical constituents of *Saxifraga stolonifera*. Zhongguo Shiyang Fangjixue Zazhi (= Chinese Journal of Experimental Medical Formulae) 2012;18:124–6.
- [142] Schütz B, Orjala J, Sticher O, Rali T. Dammarane triterpenes from the leaves of *Securinega melanthoides*. J Nat Prod 1998;61:96–8.
- [143] Sanogo R, Vassallo A, Malafronte N, Imperato S, Russo A, Piazz FD. New phenolic glycosides from *Securinega virosa* and their antioxidant activity. Nat Prod Commun 2009;4:1645–1650.
- [144] Carruthers WR, Hay JE, Haynes LJ. Isolation of bergenin from *Shorea leprosula*; identity of vakerin and bergenin. Chem Ind (London, UK) 1957;76–7.
- [145] Mukherjee H, Ojha D, Bharitkar YP, Ghosh S, Mondal S, Kaity S et al. Evaluation of the wound healing activity of *Shorea robusta*, an Indian ethnomedicine, and its isolated constituent(s) in topical formulation, J Ethnopharmacol 2013;149:335–343.
- [146] Varshney VK, Dayal R. Chemical constituents of *Shorea robusta*. Int J Chem Sci 2006;4:298–304.
- [147] Zhang X-H, Zhou T, Xuan L-J. A dipeptide and two glycosides from *Streptocaulon griffithii*. J Asian Nat Prod Res 2008;10:891–6.
- [148] Sridhar C, Krishnaraju AV, Subbaraju GV. Antiinflammatory constituents of *Teramnus labialis*. Indian J Pharm Sci 2006;68:111–4.
- [149] Jachak SM, Gautam R, Selvam C, Madhan H, Srivastava A, Khan T. Anti-inflammatory, cyclooxygenase inhibitory and antioxidant activities of standardized extracts of *Tridax procumbens* L. Fitoterapia 2011;82:173–7.
- [150] Zhang T, Li B, Chen L, Li J-J, Liu S-J, Jun J-X. A novel lactone from *Tripterospermum chinense*. Yaoxue Xuebao (= Acta Pharmaceutica Sinica) 2012;47:1517–1520.
- [151] Ito T, Abe N, Masuda Y, Nasu M, Oyama M, Sawa R et al. Two novel resveratrol derivatives from the leaves of *Vateria indica*. Helvetica Chimica Acta 2009;92:195–208.
- [152] Ito T, Masuda Y, Abe N, Oyama M, Sawa R, Takahashi Y et al. Chemical constituents in the leaves of *Vateria indica*. Chem Pharm Bull 2010;58:1369–1378.
- [153] Bhrara SC, Seshadri TR. Chemical components of *Vateria indica* seeds. Current Science 1966;35:486–7.

- [154] Ito T, Tanaka T, Iinuma M, Nakaya K-I, Takahashi Y, Sawa R et al. Two new oligostilbenes with dihydrobenzofuran from the stem bark of *Vateria indica*. *Tetrahedron* 2003;59:1255–1264.
- [155] Abe N, Ito T, Ohguchi K, Nasu M, Masuda Y, Oyama M et al. Resveratrol oligomers from *Vatica albiramis*. *J Nat Prod* 2010;73:1499–1506.
- [156] Ito T, Hara Y, Oyama M, Tanaka T, Murata J, Darnaedi D et al. Occurrence of bergenin phenylpropanoates in *Vatica bantamensis*. *Phytochem Lett* 2012;5:743–6.
- [157] Seo E-K, Chai H, Constant HL, Santisuk T, Reutrakul V, Beecher CWW et al. Resveratrol tetramers from *Vatica diospyroides*. *J Org Chem* 1999;64:6976–6983.
- [158] Song X-M, Chen G-Y, Song X-P, Han C-R, Chen S-Q, Weng S-C. Study on the chemical constituents of leaves from *Vatica mangachpoi* Blanco. *Linchan Huaxue Yu Gongye* (= Chemistry and Industry of Forest Products) 2012;32:102–6.
- [159] Mo Z, Chen G, Wang J, Wang T, Dai C, Yuan Y. The extraction technology of bergenin from leaf of *Vatica mangachpoi* Blanco. *Shipin Keji* (= Food Science and Technology) 2012;37:207–9.
- [160] Ito T, Tanaka T, Iinuma M, Iliya I, Nakaya K-I, Ali Z et al. New resveratrol oligomers in the stem bark of *Vatica pauciflora*. *Tetrahedron* 2003;59:5347–5363.
- [161] Khosa RL, Wahi AK, Mohan Y, Ray AB. Isolation of bergenin from the roots of *Viburnum nervosum*. *Indian J Pharm Sci* 1978;67–9.
- [162] Tiwari BK, Khosa RL. Studies on *Viburnum nervosum* Hook: Chemistry and spectroscopy of bergenin and its derivatives. *International J Pharmaceutical Sciences and Research* 2012;3:1361–3 (<http://www.oalib.com/paper/2738963>).
- [163] Kalidhar SB, Parthasarathy MR, Sharma P. Norbergenin, a new C-glycoside from *Woodfordia fruticosa* Kurtz. *Indian J Chem* 1981;20B:720–1.
- [164] Patel DK, Patel K, Kumar R, Gadewar M, Tahilyani V. Pharmacological and analytical aspects of bergenin: A concise report. *Asian Pacific Journal of Tropical Biomedicine* 2012;163–7.
- [165] Chopra RN, Maya SL, Chopra IC. Glossary of Indian Medicinal Plants. Council of Scientific and Industrial Research, New Delhi; 1956, p. 156.
- [166] Li Y, Liu J, Song G, Li K, Zhang K, Ye B. Sensitive voltammetric sensor for bergenin based on poly(L-lysine)/grapheme modified glassy carbon electrode. *Anal Methods* 2013;5:3895–3902.
- [167] Ghimire SK, Sapkota IB, Oli BR, Parajuli RR. Non-Timber Forest Products of Nepal Himalaya (Database of some important species found in the mountain protected areas and surrounding regions). WWF Nepal, Kathmandu, Nepal; 2008.
- [168] Chinese Pharmacopoeia Commission. Pharmacopoeia of the People's Republic of China. vol 1. China Medical Science Press; 2010.
- [169] Khare CP, editor. Indian Medicinal Plants: An Illustrative Dictionary. Springer-Verlag Berlin: Heidelberg; 2007.
- [170] Shukla YN, Bhakuni RS, Singh DP, Jain SP. The genus *Astilbe* – A review on its chemical and pharmacological research. *Journal of Medicinal and Aromatic Plant Sciences* 2004;26:324–331.
- [171] Kobayashi H, De Mejía E. The genus *Ardisia*: A novel source of health-promoting compounds and phytopharmaceuticals. *J Ethnopharmacol* 2005;96:347–354.
- [172] Rastogi S, Rawat AKS. A comprehensive review on bergenin, a potential hepatoprotective and antioxidative phytoconstituent. *Herba Polonica* 2008;54:66–79.
- [173] Singh N, Gupta AK, Juyal V. A review on *Bergenia Ligulata* Wall. *International Journal of Chemical and Analytical Science* 2010;1:71–3 (<http://ijcas.info/index.php/ijca/article/view/2155>).
- [174] Zhang Y, Liao C, Liu X, Li J, Fang S, Li Y et al. Biological advances in *Bergenia* genus plant. *African J Biotechnol* 2011;10:8166–9.
- [175] Sharma J, Varmal R. A review on endangered plant of *Mallotus philippensis* (Lam.) M. Arg. *Pharmacologyonline* 2011;3:1256–1265 (pharmacologyonline.silae.it/files/newsletter/2011/vol3/124.sharma.pdf)
- [176] Ruby Km, Chauhan R, Sharma S, Dwivedi J. Polypharmacological activities of *Bergenia* species. *Int J Pharm Sci Rev Res* 2012;13:100–110 (globalresearchonline.net/journalcontents/v13-1/018.pdf).
- [177] Chauhan R, Ruby Km, Dwivedi J. Himalayan *Bergenia* a comprehensive review. *Int J Pharm Sci Rev Res* 2012;14:139–141 (globalresearchonline.net/journalcontents/v14-2/24.pdf).

- [178] Chauhan R, Ruby Km, Dwivedi J. *Bergenia ciliata* mine of medicinal properties: A review. Int J Pharm Sci Rev Res 2012;15:20–3 (globalresearchonline.net/journalcontents/v15-2/04.pdf).
- [179] Chauhan R, Ruby Km, Dwivedi J. Secondary metabolites found in *Bergenia* species: A compendious review. Int J Pharm Pharm Sci 2013;5:9–16 (www.ijppsjournal.com/Vol5Issue1/6027.pdf).
- [180] Chen P, Lei J, Xu X, Yang J. Chemical constituents and antibacterial activity contained in *Caesalpinia millettii*. Zhongguo Zhongyao Zazhi (= China Journal of Chinese Materia Medica) 2012;37:2105–7.
- [181] Piegen X. Traditional experience of Chinese herb medicine. Its application in drug research and new drug searching. In: Beal JL, Reinhard E, editors. Natural Products as Medicinal Agents. Hippokrates Verlag: Stuttgart, Germany; 1980, p. 351–394.
- [182] Xie JX, Wang L, Liu CX, Zhang DY. The identification total synthesis of aichasu, an antitussive agent. Acta Pharmacol Sin 1981;6:425–8.
- [183] Prithiviraj B, Singh UP, Manickam M, Srivastava JS, Ray AB. Antifungal activity of bergenin, a constituent of *Flueggea microcarpa*. Plant Pathology 1997;46:224–8.
- [184] Hikino H, Kiso Y, Hatano T, Yoshida T, Okuda T. Antihepatotoxic actions of tannins. J Ethnopharm 1985;14:19–29.
- [185] Lim H-K, Kim H-S, Chung M-W, Kim YC. Protective effects of bergenin, the major constituent of *Mallotus japonicus*, on D-galactosamine-intoxicated rat hepatocytes. J Ethnopharmacol 2000;70:69–72.
- [186] Lim H-K, Kim H-S, Choi H-S, Oh S, Choi J. Hepatoprotective effects of bergenin, a major constituent of *Mallotus japonicus*, on carbon tetrachloride-intoxicated rats. J Ethnopharmacol 2000;72:469–474.
- [187] Newell AMB, Yousef GG, Lila MA, Ramírez-Mares MV, De Meijia EG. Comparative in vitro bioactivities of tea extracts from six species of *Ardisia* and their effect on growth inhibition of HepG2 cells. J Ethnopharmacol 2010;130:536–544.
- [188] Singh U, Barik A, Priyadarsini KI. Reactions of hydroxyl radical with bergenin, a natural poly phenol studies by pulse radiolysis. Bioorg Med Chem 2009;17:6008–6014.
- [189] Yokozawa T, Chen C-P, Dong E, Tanaka T, Nonaka G-I, Nishioka I. Study on the inhibitory effect of tannins and flavonoids against the 1,1-diphenyl-2-picrylhydrazyl radical. Biochemical Pharmacology 1998;56:213–222.
- [190] Okada T, Suzuki T, Hasobe S, Kisara K. Studies on bergenin (report I) – Antiulcerogenic activities of bergenin. Folia Pharmacologica Japonica 1973;69:369–378 (https://www.jstage.jst.go.jp/article/fpj1944/69/.../_pdf).
- [191] Abe K, Sakai K, Uchida M. Effects of bergenin on experimental ulcers – Prevention of stress induced ulcers in rats. General Pharmacology: The Vascular System 1980;11:361–8.
- [192] Goel RK, Maiti RN, Manickam M, Ray AB. Antiulcer activity of naturally occurring pyranocoumarin and isocoumarins and their effect on prostanoid synthesis using human colonic mucosa. Indian J Experimental Biology 1997;35:1080–3.
- [193] Satish H, Umashanker DC. Comparative study of methanolic extract of *Bergenia ligulata* Yeo. with isolated constituent bergenin in urolithiatic rats. Biomed 2006;1:80–6.
- [194] Dharmender R, Madhavi T, Reena A, Sheetal A. Simultaneous quantification of bergenin, (+)-catechin, gallicin and gallic acid; and quantification of β -sitosterol using HPTLC from *Bergenia ciliata* (Haw.) Sternb. *Forma ligulata* Yeo (Pasanbheda). Pharm Anal Acta 2010;1:104 (<http://omicsonline.org/2153-2435/2153-2435-1-104.php>).
- [195] Dhalwal K, Shinde VM, Biradar YS, Mahadik KR. Simultaneous quantification of bergenin, catechin and gallic acid from *Bergenia ciliata* and *Bergenia ligulata* by using thin-layer chromatography. J Food Composition and Analysis 2008;21:496–500.
- [196] Chauhan SK, Singh B, Agrawal S. Simultaneous determination of bergenin and gallic acid in *Bergenia ligulata* Wall by high-performance thin layer chromatography. J AOAC Int 2000;83:1480–3.
- [197] Srivastava S, Rawat AKS. Simultaneous determination of bergenin and gallic acid in different *Bergenia* species. Journal of Planar Chromatography – Modern TLC 2007;20:275–7.
- [198] Haribabu K, Ajitha M, Ramesh B, Babu KS, Rao JM. Quantification of bergenin from *Mallotus philippinensis* by HPTLC-MS and study on different extraction methods. Journal of Planar Chromatography – Modern TLC 2012;25:445–9.


- [199] Singh DP, Srivastava SK, Govindarajan R, Rawat AKS. High-performance liquid chromatographic determination of bergenin in different *Bergenia* species. *Acta Chromatographica* 2007;19:246–252.
- [200] Jiang H-J, Guo F-G, Zhang L-M, Chen Y-P, Yang S-C. Comparison of bergenin contents of *Bergenia purpurascens* among different regions in Yunan Province. *Journal of Yunnan Agricultural University (Natural Science)* 2010;25:895–8 (<http://xb.ynau.edu.cn/CN/abstract/abstract9560.shtml>).
- [201] Ji L-J. Bergenin HPLC determination of two species of *Bergenia* growing in Tibet. *Acta Botanica Boreali-Occidentalia Sinica* 2005;397–9.
- [202] Cheng B, Li Y-X, Quan Q-M, Xiao H, Gong X-Y. Comparison of bergenin contents of *Bergenia emeiensis* in different altitude in E mei. *Journal of China West Normal University (Natural Science)* 2011;317–320.
- [203] Raj MK, Balachandran C, Duraipandiyan V, Agastian P, Ignacimuthu S, Vijaykumar A. Isolation of terrestribisamide from *Peltophorum pterocarpum* (DC.) Baker ex. K. Heyne and its antimicrobial, antioxidant, and cytotoxic activities. *Med Chem Res* 2013;22:3823–3830.
- [204] Zhang Z-P, Pan Z-G, Zhang J-P. The separation and detection of bergenin from herba *Ardisiae japonicae*. *Shipin Gongye (Shanghai, China)* 2012;33:131–3.
- [205] Veerapur VP, Prabhakar KR, Thippeswamy BS, Bansal P, Srinivasan KK, Unnikrishnan MK. Antidiabetic effect of *Ficus racemosa* Linn. stem bark in high-fat diet and low-dose streptozotocin-induced type-2 diabetic rats: A mechanistic study. *Food Chemistry* 2012;132:186–193.
- [206] Ahmed F, Urooj A. Cardioprotective activity of standardized extract of *Ficus racemosa* stem bark against doxorubicin-induced toxicity. *Pharmaceutical Biol* 2012;50:468–473.
- [207] Shi Y-B, Shi Y-P, Meng Q-G. Determination and pharmacokinetics study of bergenin in rat plasma by RP-HPLC method. *Biomedical Chromatography* 2006;20:1065–1070.
- [208] Qin X, Zhou D, Zhang Z-R, Huang Y. Determination of bergenin in rat plasma by high-performance liquid chromatography. *Pharmazie* 2007;62:323–6.
- [209] Shi Y-B, Shi Y-P, Zhang X-Y, Zhao Q-Y, Ni J-M. Determination of bergenin in rat urine, feces and tissues by RP-HPLC method. *Journal of Chinese Pharmaceutical Sciences* 2009;18:49–54.
- [210] Yu W, Wang Y, Zhang Y, Zhang D, Lan J, Liu Z et al. Quantitation of bergenin in human plasma by liquid chromatography/tandem mass spectrometry. *J Chromatography B* 2009;877:33–6.
- [211] Wang J, Wang B-J, Wei C-M, Yuan G-Y, Zhang R, Liu H et al. Determination of bergenin in human plasma after oral administration by HPLC-MS/MS method and its pharmacokinetic study. *Biomed Chromatogr* 2009;23:199–203.
- [212] Zhang Y, Dong L, Li Y, Li J, Chen X. Characterization of interaction between bergenin and human serum albumin in membrane mimetic environments. *J Fluoresc* 2008;18:661–670.
- [213] Qin X, Yang Y, Fan T-T, Gong T, Zhang X-N, Huang Y. Preparation, characterization and in vivo evaluation of bergenin-phospholipid complex. *Acta Pharmacologica Sinica* 2010;31:127–136.
- [214] Chen J, Zhang J, Zhuang Q, Zhang S, Lin X. Electrochemical study of bergenin on a poly(4-(2-pyridylazo)-resorcinol) modified glassy carbon electrode and its determination in tablets and urine. *Talanta* 2007;72:1805–1810.
- [215] Zhuang Q, Chen J, Chen J, Lin X. Electrocatalytic properties of bergenin on a multi-wall carbon nanotubes modified carbon paste electrode and its determination in tablets. *Sensors and Actuators B* 2008;128:500–6.
- [216] Feng W-S, Li Z, Zheng X-K, Li Y-J, Su F-Y, Zhang Y-L. Chemical constituents of *Saxifraga stolonifera* (L.) Meeb. *Yao Xue Xue Bao (= Acta Pharmaceutica Sinica)* 2010;45:742–6.
- [217] Nyunt KS, Elkhateeb A, Tosa Y, Nabata K, Katakura K, Matsuura H. Isolation of antitrypanosomal compounds from *Vitis repens*, a medicinal plant of Myanmar. *Nat Prod Commun* 2012;7:609–610.
- [218] Vaishali AS, Vikas MD, Krishnapriya M, Sanjeevani G. Identification of potential antioxidants by in-vitro activity guided fractionation of *Bergenia ligulata*. *Pharmacognosy Magazine* 2008;4:79–84.
- [219] Chandraredy UD, Chawla AS, Mundkinajeddu D, Maurya R, Handa SS. Paashaanolactone from *Bergenia ligulata*. *Phytochemistry* 1998;47:907–9.
- [220] Arfan M, Amin H, Khan N, Khan I, Saeed M, Khan MA et al. Analgesic and anti-inflammatory activities of 11-O-galloylbergenin. *J Ethnopharmacol* 2010;131:502–4.

- [221] Janar J, Fang L, Chin CP, Kaneda T, Hirasawa Y, Shahmanovna B. A new galloylbergenin from *Bergenia crassifolia* with anti-lipid droplet accumulation activity. *Heterocycles* 2012;86:1591–5.
- [222] Lee YY, Jang DS, Jin JL, Yun-Choi HS. Antiplatelet aggregating and antioxidative activities of 11-O-(4'-O-methylgalloyl)-bergenin, a new compound isolated from *Craddula* cv. 'Himaturi'. *Planta Medica* 2005;71:776–7.
- [223] Zuo G-Y, Li Z-Q, Chen L-R, He H-P, Xu X-J. Gallic acid esters of bergenin from *Saxifraga melanocentra* (Saxifragaceae) and their inhibition against HCV NS3 protease. *Acta Botanica Yunnanica* 2007;29:486–8.
- [224] Takahashi H, Kosaka M, Watanabe Y, Nakade K, Fukuyama Y. Synthesis and neuroprotective activity of bergenin derivatives with antioxidant activity, *Bioorg Med Chem* 2003;11:1781–8.
- [225] Jung J-C, Lim E, Kim SH, Kim NS, Jung M, Oh S. Practical synthesis and biological evaluation of bergenin analogs. *Chem Biol Drug Res* 2011;78:725–9.
- [226] Lim H-K, Kim H-S, Choi H-S, Oh S, Jang C-G, Choi J et al. Effects of acetylbergenin against D-galactosamine-induced hepatotoxicity in rats. *Pharmacological Research* 2000;42:471–4.
- [227] Mozhaev VV, Budde CL, Rich JO, Usyatinsky A Ya, Michels PC, Khmelnsky YL et al. Regioselective enzymatic acylation as a tool for producing solution-phase combinatorial libraries. *Tetrahedron* 1998;54:3971–3982.
- [228] Kashima Y, Miyazawa M. Synthesis, antioxidant capacity, and structure-activity relationships of tri-O-methylnorbergenin analogues on tyrosinase inhibition. *Bioorg Med Chem Lett* 2013;23:6580–4.
- [229] Wang D, Zhu H-T, Zhang Y-J, Yang C-R. A carbon-carbon-coupled dimeric bergenin derivative biotransformed by *Pleurotus ostreatus*. *Bioorg Med Chem Lett* 2005;15:4073–5.
- [230] Minamikawa T, Yoshida S, Hasegawa M, Komagata K, Kato K. Microbial degradation of bergenin, a phenolic C-glucoside. *Agric Biol Chem* 1972;36:773–8.
- [231] Hattori M, Shu Y-Z, Tomimori T, Kobashi K, Namba T. A bacterial cleavage of the C-glucosyl bond of mangiferin and bergenin. *Phytochemistry* 1989;28:1289–1290.
- [232] Schmidt RR, Effenberger G. Aryl-C-glucosides from O- α -D-glucosyl trichloroacetimidate – Structure of bergenin derivatives. *Carbohydrate Res* 1987;171:59–79.
- [233] Hua X-G, Mague JT, Li C-J. Model studies of (+)-bergenin: a convenient formation of aryl δ -lactones. *Tetrahedron Lett* 1998;39:6837–6840.
- [234] Frick W, Schmidt RR. Synthesis of bergenin-type C-glucosylarenes. *Carbohydrate Res* 1991;209:101–7.
- [235] Rousseau C, Martin OR. Synthesis of bergenin-related natural products by way of an intramolecular C-glycosylation reaction. *Tetrahedron Asymmetry* 2000;11:409–412.
- [236] Herzner H, Palmacci ER, Seeberger PH. Short total synthesis of 8,10-di-O-methylbergenin. *Org Lett* 2002;4:2965–7.
- [237] Sakamaki S, Kawanishi E, Nomura S, Ishikawa T. Aryl- β -C-glucosidation using glucal boronate: Application to the synthesis of tri-O-methylnorbergenin. *Tetrahedron* 2012;68:5744–5753.
- [238] Parkan K, Pohl R, Kotora M. Cross-coupling reaction of saccharide-based alkenyl boronic acids with aryl halides: The synthesis of bergenin. *Chem Eur J* 2014;20:4414–9.
- [239] Farnsworth NR, Akerele O, Binanggal AS, Soejarto DD, Guo Z. Medicinal Plants in Therapy. *Bulletin of the World Health Organization* 1985;63:965–981.
- [240] Sharma HK, Chhangte L, Dolui AK. Traditional medicinal plants in Mizoram, India. *Fitoterapia* 2001;72:146–161.
- [241] Ballabh B, Chaurasia OP, Ahmed Z, Singh SB. Traditional medicinal plants of cold desert Ladakh – Used against kidney and urinary disorders. *J Ethnopharmacol* 2008;118:331–9.

Graphical Abstract

Diversity, pharmacology and synthesis of bergenin and its derivatives: Potential materials for therapeutic usages

Gan B. Bajracharya



List of compounds discussed in the article and are entered in the NCBI PubChem Compound database:

Bergenin (PubChem CID: 66065)

Norbergenin (PubChem CID: 73192)

4-O-Galloylbergenin (PubChem CID: 14464332)

11-O-Galloylbergenin (PubChem CID: 14464334)

11-O-Syringylbergenin (PubChem CID: 195481)